

STATION 144A
TO 145A.

- Davila* (?) *umbonata*, n.sp. (100 and 150 fathoms); obtained also at Station 149, 20 to 60 fathoms.
- Cryptodon marionensis*, n.sp. (100 to 150 fathoms); obtained at no other locality.
- Kellia nuculina*, Martens. (50 to 150 fathoms); obtained also at Station 149, 20 to 120 fathoms.
- Astarte magellanica*, Smith. (100 and 150 fathoms); obtained at no other locality by the Challenger. Recorded from Magellan Strait.
- Limopsis marionensis*, n.sp. (100 to 150 fathoms); obtained at no other locality.
- Mytilus meridionalis*, n.sp. (100 to 150 fathoms); obtained also at Station 150 150 fathoms.
- Crenella marionensis*, n.sp. (140 fathoms); obtained at no other locality.
- Modiolarca trapezina* (Lamarck). (50 to 100 fathoms); obtained also at Stations 149, 311, and 315, surface to 245 fathoms.
- Dacrydium meridionalis*, n.sp. (100 to 150 fathoms); obtained at no other locality.
- Lima* (*Limatula*) *pygmæa*, Philippi. (50 to 150 fathoms); obtained also at Station 149, 28 to 60 fathoms. Recorded from South Patagonian region.
- Pecten distinctus*, n.sp. (100 fathoms); obtained at no other locality.
- „ *aviculoides*, n.sp. (100 to 150 fathoms); obtained at no other locality.

SCAPHOPODA and GASTEROPODA (Watson, Zool. pt. 42).

- Dentalium entalis*, Linné, var. *orthrum*, nov. (140 fathoms); obtained also at Stations II. and 75.
- Emarginula* sp. (?). (50 to 150 fathoms).
- Puncturella noachina* (Linné), var. *princeps*, Mighels. (69, 140, and 310 fathoms); obtained also at Stations 149 and 312, 10 to 60 fathoms. A widely-distributed species. Fossil—Miocene (?) of Sicily, Pliocene onwards of Europe, Labrador, and Nova Zembla.
- Trochus* (*Photinula*) *expansus* (Sowerby). (50 to 140 fathoms); obtained also at Station 149 and Falkland Islands, surface to 60 fathoms. Recorded from Magellan Strait.
- Scissurella crispata*, Fleming. (140 fathoms); obtained also at Station 24.
- Homalogyra atomus* (Philippi). (140 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil—later glacial beds of Norway and Scotland.
- Cyclostrema* (?) sp. (50 to 150 fathoms).
- Trophon declinans*, n.sp. (69 fathoms); obtained also at Station 150, 150 fathoms.
- Fusus* (*Neptunea*) *edwardiensis*, n.sp. (140 fathoms); obtained at no other locality.