

- Pleurotoma (Thesbia) translucida*, n.sp. (140 fathoms); obtained also at Station 149, 25 and 28 fathoms. STATIONS 144A TO 145A.
- Triton (Lagena) magellanicus* (Chemnitz). (69 fathoms); obtained also at Station 309 (?), 40 fathoms. Recorded from Magellan Strait and Cape Horn.
- „ ( „ ) n.sp. One young specimen (140 fathoms).
- Natica fertilis*, n.sp. (50 to 140 fathoms); obtained also at Stations 149 and 150, 60 and 150 fathoms.
- Turritella austrina*, n.sp. (50 to 100 fathoms); obtained also at Station 149, 28 fathoms.
- Odostomia rissoïdes*, Hanley. One specimen (50 to 140 fathoms); obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil—Upper Pliocene onwards.
- Eulima amblicia*, n.sp. One specimen and fragment (50 to 150 fathoms); obtained at no other locality.
- Cerithium* sp. (?). (100 fathoms).
- Alaba (Diala) limnæiformis*, n.sp. (50 to 150 fathoms); obtained at no other locality.
- Jeffreysia edwardiensis*, n.sp. (310 fathoms); obtained at no other locality.
- Rissoa (Ceratia) transenna*, n.sp. (140 fathoms); obtained at no other locality.
- „ (*Setia*) *marionensis*, n.sp. (50 to 140 fathoms); obtained at no other locality.
- „ ( „ ) *principis*, n.sp. (140 fathoms); obtained also at Kerguelen, shore.
- „ ( „ ) *edwardiensis*, n.sp. (50 to 150 fathoms); obtained at no other locality.
- Lamellaria* sp. (?). (50 to 75 fathoms).

## POLYPLACOPHORA (Haddon, Zool. pt. 43).

*Lepidopleurus dorsuosus*, n.sp. (310 fathoms); obtained at no other locality.

## MARSENIADÆ (Bergh, Zool. pt. 41).

*Marseniopsis murrayi*, n.g., n.sp. One specimen (50 to 75 fathoms); obtained at no other locality.

## POLYZOA (Busk, Zool. pts. 30 and 50; Waters, pt. 79).

*Menipea flagellifera*, n.sp. (50 to 75 fathoms); obtained also at Stations 149, 313, and 314, 20 to 127 fathoms.

„ *marionensis*, n.sp. (50 to 75 fathoms); obtained also at Station 142.

*Caberea darwinii*, Busk. (50 to 150 fathoms); for distribution see Station 135.