

KERGUELEN.

Psolus ephippifer, Wyville Thomson, n.sp. One specimen (20 to 60 fathoms); obtained also at Stations 145, 150, and 151.

Thyone recurvata, n.sp. One specimen (10 to 100 fathoms); obtained at no other locality.

ENTOZOA (Linstow, Zool. pt. 71).

Ascaris simplex, Rudolphi. Thirteen specimens from stomach of seal (*Otaria jubata*).

„ *spiculigera*, Rudolphi. Thirty-six specimens from stomach of shag (*Phalacrocorax verrucosus*).

NEMERTEA (Hubrecht, Zool. pt. 54).

Drepanophorus serraticollis, Hubrecht (?). One fragmentary specimen (depth not given); obtained also at Station 162, 38 to 40 fathoms. Recorded from Mediterranean.

Amphiporus moseleyi, n.sp. Several specimens (shallow water to 20 fathoms); obtained at no other locality.

„ *marioni*, n.sp. One specimen (120 fathoms); obtained also at Station 145.

Cerebratulus corrugatus (M'Intosh). Numerous specimens (25 to 120 fathoms); obtained also at Station 151, 75 fathoms.

„ sp. (?) (*medullatus*, n.sp. ?). One fragmentary specimen (depth not given).

GEPHYREA (Selenka, Zool. pt. 36).

Phascolosoma pudicum, n.sp. Numerous specimens (10 to 127 fathoms); obtained at no other locality.

ANNELIDA (M'Intosh, Zool. pt. 34).

Lætmonice producta, Grube. Numerous specimens (20 to 120 fathoms); obtained also at Station 151, 75 fathoms. Recorded from Kerguelen.

Lagisca antarctica, n.sp. Several specimens (127 fathoms); obtained also at Station 145.

„ *magellanica*, n.sp. (127 fathoms); obtained also at Station 308, 175 fathoms.

Evarne kerguelensis, n.sp. Several specimens (30 to 127 fathoms); obtained at no other locality.

Eupolynoë mollis, M'Intosh. One fragmentary specimen (100 fathoms); obtained at no other locality by the Challenger.