

- Tritata kergueleni*, n.sp. Three specimens (28 to 127 fathoms); obtained at no other locality. KERGUELEN.
- Rhachotropis kergueleni*, n.sp. Two specimens (depth not given); obtained at no other locality.
- Eusirus longipes*, Boeck. One specimen (depth not given); obtained also at Station 150, 150 fathoms. Recorded from North Atlantic.
- Liljeborgia consanguinea*, n.sp. One specimen (20 fathoms); obtained also at Station 151, 75 fathoms.
- Photis macrocarpus*, n.sp. Several specimens (depth not given); obtained at no other locality.
- Aora kergueleni*, n.sp. Three specimens (30 to 38 fathoms); obtained at no other locality.
- „ *trichobostrychus*, n.sp. Two specimens (depth not given); obtained at no other locality.
- Autonoe kergueleni*, n.sp. Two specimens (127 fathoms); obtained at no other locality.
- Gammaropsis exsertipes*, n.sp. Three specimens (depth not given); obtained at no other locality.
- Amphithoë kergueleni*, n.sp. One specimen (depth not given); obtained at no other locality.
- Podocerus falcatus* (Montagu). One specimen (30 fathoms); taken also from screw after leaving the Cape.
- Cerapus sismithi*, n.sp. Three specimens (120 fathoms); obtained at no other locality.
- Haplocheira plumosa*, n.sp. Two specimens (shallow water to 127 fathoms); obtained at no other locality.
- Platophium danaë*, n.sp. Nine specimens (127 fathoms); obtained at no other locality.
- Neohela serrata*, n.sp. Two specimens (127 fathoms); obtained at no other locality.
- Kerguelenia compacta*, n.g., n.sp. Two specimens (127 fathoms); obtained at no other locality. Only species of the genus.
- Dodecas elongata*, n.g., n.sp. Several specimens (95 to 110 fathoms); obtained at no other locality. Only species of the genus.
- Protellopsis kergueleni*, n.g., n.sp. Two specimens (30 fathoms); obtained at no other locality. Only species of the genus.
- Euthemisto thomsoni*, n.n. One specimen (25 fathoms); obtained at the surface on several occasions in the Antarctic.