

## KERGUELEN.

- Onchopora sinclairii*, Busk. (28 fathoms); obtained also at Stations 150, 151, and 157, 75 to 1950 fathoms. Recorded from New Zealand, Australia, and Marion Island. [Waters calls it *Calwellia sinclairii* (Busk)].
- Chorizopora hyalina*, var. *bougainvillei*, d'Orbigny. (28 fathoms); obtained also at Stations 145 and 315.
- Smittia marionensis*, Busk. (28 fathoms); obtained also at Station 145.
- Cellepora eatonensis*, n.sp. (28 to 127 fathoms); obtained also at Stations 303 and 315, 1325 and 5 to 12 fathoms.
- Crisia eburnea* (Linné), var. *laxa*, nov. (28 to 105 fathoms); obtained at no other locality by the Challenger. Recorded from Arctic and North Atlantic.
- Idmonea atlantica*, Forbes. (30 fathoms); obtained also at Stations 49, 135, and the Cape.

## BRACHIOPODA (Davidson, Zool. pt. 1).

- Waldheimia kerguelenensis*, n.sp. Numerous specimens (20 to 60 fathoms); obtained also at Stations 145 and 150.
- Terebratella dorsata* (Gmelin). Three specimens (20 to 30 fathoms); obtained at no other locality by the Challenger. Recorded from coast of Chili to Magellan Strait.

## TUNICATA (Herdman, Zool. pts. 17 and 38).

- Eugyra kerguelenensis*, n.sp. Three specimens (10 to 100 fathoms); obtained at no other locality.
- Styela lactea*, n.sp. Three specimens (10 to 100 fathoms); obtained at no other locality.
- Ascidia challengerii*, n.sp. Thirteen specimens (10 to 60 fathoms); obtained at no other locality.
- „ *vasculosa*, n.sp. One test (28 fathoms); obtained at no other locality.
- „ *translucida*, n.sp. Three specimens (28 fathoms); obtained at no other locality.
- „ *despecta*, n.sp. One specimen (10 to 100 fathoms); obtained at no other locality.
- Colella pedunculata* (Quoy and Gaimard). Four specimens (10 to 60 fathoms); obtained also at Stations 151, 313, 314, and 315, 12 to 75 fathoms. Recorded from Australia and Magellan Strait.