

- Colella quoyi*, n.g., n.sp. Three specimens (25 fathoms); obtained at no other locality. KERGUELEN.
- „ *concreta*, n.g., n.sp. Several specimens (10 to 60 fathoms); obtained at no other locality.
- (?) *pyriformis*, n.sp. Ten specimens without Ascidiozooids, therefore the genus is doubtful (attached to *Macrocystis pyrifer*); obtained at no other locality.
- Tylobranchion speciosum*, n.g., n.sp. Three specimens (10 to 100 fathoms); obtained at no other locality. Only species of the genus.
- Morchellioides affinis*, n.g., n.sp. Two specimens (10 to 60 fathoms); obtained at no other locality. Only species of the genus.
- Morchellium giardi*, n.sp. Four specimens (20 to 60 fathoms); obtained at no other locality.
- Polyclinum pyriformis*, n.sp. Two specimens (10 to 60 fathoms); obtained at no other locality.
- „ *minutum*, n.sp. One specimen (20 to 60 fathoms); obtained at no other locality.
- Aplidium fuscum*, n.sp. One specimen (20 to 60 fathoms); obtained at no other locality.
- „ *leucophæum*, n.sp. One specimen (10 to 60 fathoms); obtained at no other locality.
- „ *fumigatum*, n.sp. Four specimens (10 to 100 fathoms); obtained also at Philippines.
- Amaroucium variabile*, n.sp. Numerous specimens (10 to 100 fathoms); obtained at no other locality.
- „ „ var. *tenerum*, nov. Five specimens (10 to 60 fathoms); obtained at no other locality.
- „ *globosum*, n.sp. Two specimens (10 to 60 fathoms); obtained at no other locality.
- „ *complanatum*, n.sp. Five specimens (50 to 120 fathoms); obtained at no other locality.
- „ *nigrum*, n.sp. One specimen (28 fathoms); obtained at no other locality.
- Psammaplidium retiforme*, n.g., n.sp. One specimen and fragments (50 to 120 fathoms); obtained at no other locality.
- Leptoclinum subflavum*, n.sp. One specimen (28 fathoms); obtained at no other locality.
- „ *rubicundum*, n.sp. One specimen (20 to 60 fathoms); obtained at no other locality.