DEEP-SEA MEDUSÆ (Haeckel, Zool. pt. 12).

STATION 153.

Thamnostylus dinema, n.g., n.sp. Obtained at no other locality. Only species of the genus. [Reported in error from 120 fathoms].

ASTEROIDEA (Sladen, Zool. pt. 51).

Pararchaster antarcticus, n.g., n.sp. Two specimens; obtained at no other locality.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiacantha cosmica, n.sp. For distribution see Station 122.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Goniocidaris canaliculata, Agassiz. For distribution see Station 147.

Cystechinus vesica, n.g., n.sp. Obtained also at Stations 298 and 299, 2225 and 2160 fathoms.

CIRRIPEDIA (Hoek, Zool. pt. 25).

Scalpellum antarcticum, n.sp. One specimen; obtained at no other locality.

ISOPODA (Beddard, Zool. pt. 48).

Iolanthe acanthonotus, n.g., n.sp. One specimen; obtained at no other locality.
Only species of the genus.

Arcturus furcatus, Studer. One specimen; obtained also at Stations 149 and 151.

" glacialis, n.sp. One specimen; obtained at no other locality.

Schizopoda (Sars, Zool. pt. 37).

Pseudomma sarsii, Willemoes-Suhm, n.sp. One specimen; obtained also at Station 149.

Pycnogonida (Hoek, Zool. pt. 10).

Nymphon meridionale, n.sp. One specimen; obtained at no other locality.

POLYZOA (Busk, Zool. pt. 30).

Farciminaria magna, n.sp. Obtained also at Stations 323 (var. armata) and 325, 1900 and 2650 fathoms.

In addition to the foregoing, the following are recorded in the Station-book:—Silicisponge, Farrea fecunda, Aphroditacean, and Ascidian.

Excluding Protozoa, over 20 specimens of invertebrates were obtained at this Station, belonging to about 18 species, of which 11 are new to science, including representatives of 5 new genera; 7 of the new species and 3 new genera were not obtained elsewhere.