

ASTEROIDEA (Sladen, Zool. pt. 51).

STATION 158.

- Hymenaster nobilis*, Wyville Thomson, n.sp. Obtained at no other locality.
Recorded subsequently from Indian Ocean ("Investigator").
- „ *formosus*, n.sp. Obtained at no other locality.
- „ *sacculatus*, n.sp. Obtained at no other locality.
- „ *cælatus*, n.sp. Obtained at no other locality.
- „ *crucifer*, n.sp. Obtained at no other locality.

OPHIUROIDEA (Lyman, Zool. pt. 14).

- Ophioglypha loveni*, n.sp. (?). Obtained also at Stations 146, 147, 157, and 160.
- „ *minuta*, n.sp. Obtained also at Station 146.
- Ophiacantha cosmica*, n.sp. For distribution see Station 122.

ECHINOIDEA (Agassiz, Zool. pt. 9).

- Urechinus naresianus*, n.g., n.sp. Obtained also at Stations 146, 147, and 302.
Only species of the genus.
- Cystechinus wyvillii*, n.g., n.sp. Obtained also at Stations 146, 147, 296, 299,
and 300.

HOLOTHURIOIDEA (Théel, Zool. pts. 13 and 39).

- Peniagone naresi*, n.g., n.sp. One injured specimen; obtained at no other
locality.
- „ *challengeri*, n.g., n.sp. Two specimens; obtained at no other
locality.
- Lætmogone wyville-thomsoni*, n.g., n.sp. One specimen; obtained also at
Stations 147, 232, and 300.
- Benthodytes sanguinolenta*, n.g., n.sp., var. *marginata*, nov. One specimen;
the species obtained also at Stations 160, 295, and
298, 500 to 2600 fathoms.
- „ *sordida*, n.g., n.sp. One specimen; obtained also at Stations 156,
157, and 298.
- Holothuria thomsoni*, n.sp., var. *hyalina*, nov. One injured specimen; the species
obtained also at Stations 237 and 244, 1875 and
2900 fathoms.

NEMERTEA (Hubrecht, Zool. pt. 54).

- Pelagonemertes rollestoni*, Moseley, n.g., n.sp. One specimen; obtained also at
Station 236, 775 to 420 fathoms. Only species of
the genus.