

FORAMINIFERA (Brady, Zool. pt. 22).—The following species of Foraminifera were observed in the deposit from this Station (see also Murray and Renard, Deep-Sea Deposits Chall. Exp.); the pelagic species, which make up about 65 per cent. of the carbonate of lime present in the deposit, are marked thus × :—

STATION 160.  
ORGANISMS FROM  
THE DEPOSIT.

<i>Biloculina depressa</i> , d'Orbigny.	<i>Lagena fimbriata</i> , Brady.
"    "    var. <i>murrhyna</i> , Schwager.	" <i>formosa</i> , Schwager.
<i>Spiroloculina planulata</i> (Lamarek).	" <i>globosa</i> (Montagu).
" <i>tenuis</i> (Czjzek).	" <i>gracilis</i> , Williamson.
<i>Miliolina venusta</i> (Karrer).	" <i>gracillima</i> (Seguenza).
<i>Psammosphæra fusca</i> , Schulze.	" <i>hertwigiana</i> , Brady.
<i>Jaculella obtusa</i> , Brady.	" <i>lævigata</i> (Reuss).
<i>Hyperammina ramosa</i> , Brady.	" <i>lævis</i> (Montagu).
<i>Aschemonella ramuliformis</i> , Brady.	" <i>longispina</i> , Brady.
<i>Rhizammina algæformis</i> , Brady.	" <i>marginata</i> (Walker and Boys).
<i>Reophax adunca</i> , Brady.	" <i>quadralata</i> , Brady.
" <i>dentaliniformis</i> , Brady.	" <i>staphyllearia</i> , Schwager.
" <i>diffflugiformis</i> , Brady.	" <i>stelligera</i> , Brady.
" <i>distans</i> , Brady.	" <i>striata</i> (d'Orbigny).
" <i>guttifera</i> , Brady.	" <i>sulcata</i> (Walker and Jacob).
" <i>pilulifera</i> , Brady.	<i>Nodosaria communis</i> , d'Orbigny.
" <i>scorpiurus</i> , Montfort.	" <i>roemeri</i> (Neugeboren).
<i>Haplophragmium agglutinans</i> (d'Orbigny).	<i>Uvigerina aculeata</i> , d'Orbigny.
" <i>foliaceum</i> , Brady.	" <i>asperula</i> , Czjzek.
" <i>globigeriniforme</i> (Parker and Jones).	" <i>pygmaea</i> , d'Orbigny.
" <i>latidorsatum</i> (Bornemann).	× <i>Globigerina bulloides</i> , d'Orbigny.
" <i>turbinatum</i> , Brady.	× " <i>inflata</i> , d'Orbigny.
<i>Thurammina papillata</i> , Brady.	× <i>Orbulina universa</i> , d'Orbigny.
<i>Hormosina ovicula</i> , Brady.	× <i>Pullenia obliquiloculata</i> , Parker and Jones
<i>Trochammina galeata</i> , Brady.	" <i>quinqueloba</i> , Reuss.
" <i>lituiformis</i> , Brady.	" <i>sphaeroides</i> (d'Orbigny).
" <i>trullissata</i> , Brady.	<i>Sphaeroidina bulloides</i> , d'Orbigny.
<i>Textularia concava</i> (Karrer).	<i>Truncatulina akneriana</i> (d'Orbigny) (?).
<i>Verneuilina pygmaea</i> (Egger).	" <i>haidingerii</i> (d'Orbigny) (?).
<i>Gauldryina pupoides</i> , d'Orbigny.	" <i>pygmaea</i> , Hantken.
<i>Clavulina communis</i> , d'Orbigny.	" <i>ungeriana</i> (d'Orbigny).
<i>Virgulina schreibersiana</i> , Czjzek	" <i>wuellerstorfi</i> (Schwager).
" <i>squamosa</i> , d'Orbigny.	× <i>Pulvinulina crassa</i> (d'Orbigny).
" <i>subdepressa</i> , Brady.	" <i>exigua</i> , Brady.
<i>Cassidulina subglobosa</i> , Brady.	× " <i>meliniana</i> (d'Orbigny).
<i>Lagena acuticosta</i> , Reuss.	<i>Rotalia orbicularis</i> , d'Orbigny.
" <i>alveolata</i> , Brady.	<i>Nonionina depressula</i> (Walker and Jacob).
" <i>exsculpta</i> , Brady.	" <i>umbilicatula</i> (Montagu).
	<i>Polystomella striatopunctata</i> (Fichtel and Moll).

Surface Organisms.—The following species are recorded from the surface in the vicinity of this Station :—

ORGANISMS FROM  
SURFACE-NETS.