Fibularia australis, Desmoulins. (30 to 35 fathoms); obtained also at Station 164, PORT JACKSON. 950 (?) fathoms.

Peronella peroni (Agassiz). (30 to 35 fathoms); obtained also at Station 162.

Maretia planulata, Gray. (2 to 15 fathoms); obtained also at Fiji and Admiralty Islands.

Eupatagus valenciennesii, Agassiz. (30 to 35 fathoms); obtained also at Station 188, 28 fathoms.

Echinocardium australe, Gray. (6 to 15 fathoms); obtained also at Japan, 7 to 50 fathoms.

HOLOTHURIOIDEA (Théel, Zool. pt. 13).

Chirodota japonica, Marenzeller. Several specimens (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Japan.

Cucumaria mirabilis, n.sp. One specimen (6 to 15 fathoms); obtained also at Philippines, 100 fathoms.

Colochirus spinosus (Quoy and Gaimard). Eighteen specimens (6 to 7 fathoms); obtained at no other locality by the Challenger. Recorded from China and Australia.

,, australis, Ludwig. One specimen (6 to 7 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.

Phyllophorus (?) incompertus, n.sp. Two specimens (6 to 7 fathoms); obtained at no other locality.

GEPHYREA (Selenka, Zool. pt. 36).

Phymosoma japonicum, Grube. Two specimens (shallow water); obtained at no other locality by the Challenger. Recorded from Japan.

Annelida (M'Intosh, Zool. pt. 34).

Aphrodita australis, Baird. One specimen (35 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and Tasmania.

Thalenessa fimbriata, n.sp. One specimen (30 to 35 fathoms); obtained at no other locality.

Eunice aphroditois, Pallas. One specimen (2 to 10 fathoms); obtained also at Philippines. Recorded from Indian Seas.

Sabellaria (Pallasia) giardi, n.sp. One specimen (35 fathoms); obtained at no other locality.

Sabella fusca, Grube (?). One specimen (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Ceylon.