Ervilia bisculpta, Gould. (2 to 10 fathoms); obtained also at Sandwich Islands. Port Jackson. Recorded from Japan and Philippines.

- Psammobia zonalis (Lamarck). (4 to 18 fathoms); obtained at no other locality by the Challenger. Recorded from Amboina, Java, Philippines, and Tasmania.
  - modesta, Deshayes. Single valve (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- Tellina (---?) tenuilirata, Sowerby. (2 to 10 fathoms); obtained also in Torres Strait and Fiji. Recorded from Australia.
- (---?) semitorta, Sowerby. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- Donax nitidus, Deshayes. (6 to 15 fathoms); obtained also at Station 187, 6 fathoms. Recorded from Australia.
- Tapes (Paratapes) textrix (Chemnitz). (4 to 18 fathoms); obtained at no other locality by the Challenger. Recorded from Indian Ocean.
  - (Amygdala) fabagella, Deshayes. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Lake Macquarie and Tasmania.
- Venus (Chione) calophylla, Philippi. (4 to 18 fathoms); obtained also at Stations 187 and 203, 3 to 28 fathoms. Recorded from China.
  - " ( " ) jacksoni, n.sp. (4 to 18 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
  - (2 to 10 fathoms); obtained also at ( ,, ) striatissima, Sowerby. Station 162.
  - (Leukoma) australis, Sowerby. (5 to 7 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- Cytherea (Callista) disrupta, Sowerby. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- Circe scripta (Linné). (4 to 18 fathoms); obtained also in Torres Strait, 3 to 11 fathoms. Recorded from Red Sea through the Indian Ocean to Moluccas.
  - angasi, n.n. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- (6 to 7 fathoms); obtained at no other locality by Dosinia circinaria, Deshayes. the Challenger. Original locality unknown.