

PORT JACKSON.

- Cardium (Bucardium) tenuicostatum*, Lamarck. (2 to 10 fathoms); obtained also at Station 208, 18 fathoms. Recorded from Philippines, Australia, and Tasmania.
- „ („) *pulchellum*, Gray. (2 to 10 fathoms); obtained also at Stations 161 and 162.
- Chama spinosa*, Broderip. One specimen (depth not stated); obtained at no other locality by the Challenger. Recorded from Lord Hood's Island.
- Lucina ramsayi*, n.sp. (6 to 7 fathoms); obtained at no other locality.
- „ (*Loripes*?) *jacksoniensis*, n.sp. One specimen (6 to 15 fathoms); obtained at no other locality.
- Diplodonta* sp. (?). Single valve (2 to 10 fathoms).
- Kellia rotunda* (Deshayes), var. (?). One specimen (depth not stated); obtained also at Station 162.
- „ *adamsi* (Angas). Single valve (2 to 10 fathoms); obtained at no other locality by the Challenger.
- Montacuta angasi*, n.sp. Single valve (2 to 10 fathoms); obtained at no other locality.
- Cardita excavata*, Deshayes. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- „ sp. (?). Single valve (2 to 10 fathoms).
- Carditella angasi*, n.sp. (35 fathoms); obtained at no other locality.
- Trigonia lamarckii*, Gray. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- Pectunculus holosericus*, Reeve. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- „ *striatularis*, n.sp. (?) (Lamarck?), Reeve. (6 to 7 fathoms); obtained also at Station 212, 10 fathoms. Recorded from Australia.
- Limopsis brazieri*, Angas. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- Arca (Barbatia) fasciata*, Reeve. (2 to 10 fathoms); obtained at no other locality by the Challenger.
- „ (*Scapharca*) *gubernaculum*, Reeve. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Philippines.
- Mytilus hirsutus*, Lamarck. (6 to 15 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and New Zealand to China and Japan.