PORT JACKSON.	Trochus	(Clanculus) homalomphalus, Adams. (2 to 15 fathoms); obtained at
		no other locality by the Challenger. Recorded from
		Australia.
	,,	(,,) maugeri, Wood. (2 to 10 fathoms); obtained at no other
		locality by the Challenger. Recorded from Australia.
	,,	(,,) floridus, Philippi. (2 to 10 fathoms); obtained at no other
		locality by the Challenger. Recorded from Australia
		and New Ireland.
	,,	(Euchelus) baccatus (Menke). (6 to 7 fathoms); obtained at no other
	"	locality by the Challenger. Recorded from Australia,
		Solomon Islands, and Singapore.
		(Ziziphinus) decoratus, Philippi. (2 to 10 fathoms); obtained at no other
	"	locality by the Challenger. Recorded from Australia.
		(Elenchus) badius, Wood. (6 to 7 fathoms); obtained at no other locality
	,,	by the Challenger. Recorded from Australia and
		Tasmania.
		(Bankivia) fasciatus (Menke). (2 to 10 fathoms); obtained at no other
	"	
		locality by the Challenger. Recorded from Australia.
	"	(Leiopyrga) picturatus (Adams). (2 to 35 fathoms); obtained at no
		other locality by the Challenger. Recorded from
		Australia.
	,,,	(Trochocochlea) fuligineus, Adams. (2 to 10 fathoms); obtained at no
		other locality by the Challenger. Original locality
		unknown.
¥	"	(Solariella) prodictus, Fischer. (2 to 10 fathoms); obtained at no other
		locality by the Challenger. Recorded from Australia
		and Sandwich Islands.
		(,,) vitiligineus, Menke. (2 to 10 fathoms); obtained also at
		Station 186, 8 fathoms. Recorded from Australia to
		Philippines and Japan.
	,,	(,,) albugo, n.sp. (2 to 10 fathoms); obtained at no other
		locality.
	,,	, two other species undetermined (2 to 10 fathoms).
	Stomatei	la imbricata, Lamarck. (2 to 10 fathoms); obtained at no other locality
		by the Challenger. Recorded from Australia.
	,,	(Gena) nigra, Quoy and Gaimard. (2 to 10 fathoms); obtained also at
		Fernando Noronha.
	Schismo	pe carinata, n.sp. (6 to 15 fathoms); obtained also at Station 185,
		1550

155 fathoms.