- Conus (Leptoconus) jukesii, Reeve. (2 to 10 fathoms); obtained at no other Port Jackson. locality by the Challenger. Recorded from Australia.
- Triton (Simpulum) costatus (Born). (2 to 10 fathoms); obtained at no other locality by the Challenger. A widely-distributed species. Fossil—Upper Miocene of Calabria onwards.
 - ,, (Cabestana) spengleri (Chemnitz). (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and New Zealand.
- Ranella (Argobuccinum) leucostoma, Lamarck. (2 to 18 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- Cassis (Semicassis) pila, Reeve. (2 to 10 fathoms); obtained also at Stations 188 and 203, 12 to 28 fathoms. Recorded from China, Japan, and Torres Strait.
 - " sp. (?). (2 to 10 fathoms).
- Struthiolaria scutulata (Martyn). (2 to 15 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and New Zealand.
- Natica euzona, Recluz. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Philippines.
 - " sagittata, Menke. (6 to 7 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and Pacific Islands.
 - " effossa, n.sp. (30 to 35 fathoms); obtained also at Station 161.
 - ", (Neverita) conica, Lamarck. (2 to 18 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
 - ", (",) didyma, Bolten. (2 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from the whole Indian Ocean, Australia, New Guinea, and Japan.
- Crepidula aculeata (Gmelin). (2 to 10 fathoms); obtained also at Station 233, 8 to 50 fathoms. A widely-distributed species.
- Calyptræa calyptræformis (Lamarck). (2 to 15 fathoms); obtained also at Station 162.
- Turritella sinuata, Reeve. (2 to 35 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- ,, sp. (?). (2 to 10 fathoms).

 Odostomia (Parthenia) casta (Adams). (2 to 10 fathoms); obtained also at

 Station 162.