

STATIONS 164B
AND 164C.

	<i>Nodosaria raphanus</i> (Linné).	× <i>Globigerina helicina</i> , d'Orbigny.
	„ (<i>Glandulina</i>) <i>rotundata</i> , Reuss.	× „ <i>inflata</i> , d'Orbigny.
	„ <i>scalaris</i> (Batsch).	× „ <i>rubra</i> , d'Orbigny.
	„ „ var. <i>separans</i> , Brady.	× „ <i>sacculifera</i> , Brady.
	„ <i>soluta</i> , Reuss.	× <i>Orbulina universa</i> , d'Orbigny.
	„ sp. (?).	× <i>Hastigerina pelagica</i> (d'Orbigny).
	<i>Rhabdogonium tricarinatum</i> (d'Orbigny).	× <i>Pullenia obliquiloculata</i> , Parker and Jones.
	<i>Marginulina costata</i> (Batsch).	„ <i>quinqueloba</i> , Reuss.
	<i>Vaginulina spinigera</i> , Brady.	„ <i>sphaeroides</i> (d'Orbigny).
	<i>Cristellaria articulata</i> , Reuss.	<i>Sphaeroidina bulloides</i> , d'Orbigny.
	„ <i>compressa</i> , d'Orbigny.	× „ <i>dehiscens</i> , Parker and Jones.
	„ <i>convergens</i> , Bornemann.	<i>Truncatulina haidingerii</i> (d'Orbigny).
	„ <i>cultrata</i> (Montfort).	„ <i>lobatula</i> (Walker and Jacob).
	„ <i>echinata</i> (d'Orbigny).	„ <i>reticulata</i> (Czjzek).
	„ <i>gibba</i> , d'Orbigny.	„ <i>tenuimargo</i> , Brady.
	„ <i>italica</i> (Defrance).	„ <i>ungeriana</i> (d'Orbigny).
	„ <i>orbicularis</i> (d'Orbigny).	„ <i>wuellerstorfi</i> (Schwager).
	„ <i>papillosa</i> (Fichtel and Moll).	<i>Anomalina ammonoides</i> (Reuss).
	„ <i>reniformis</i> , d'Orbigny (?).	„ <i>coronata</i> , Parker and Jones.
	„ <i>rotulata</i> (Lamarck).	„ <i>grosserugosa</i> (Gümbel) (?).
	„ <i>variabilis</i> , Reuss.	„ <i>polymorpha</i> , Costa.
	„ sp. (?).	<i>Pulvinulina auricula</i> (Fichtel and Moll).
	<i>Uvigerina aculeata</i> , d'Orbigny.	× „ <i>canariensis</i> (d'Orbigny).
	„ <i>asperula</i> , Czjzek.	× „ <i>crassa</i> (d'Orbigny).
	„ „ var. <i>ampullacea</i> , Brady.	„ <i>elegans</i> (d'Orbigny).
	„ <i>pygmæa</i> , d'Orbigny.	„ <i>hauerii</i> (d'Orbigny).
	<i>Sagrina columellaris</i> , Brady.	× „ <i>menardii</i> (d'Orbigny).
	<i>Ramulina globulifera</i> , Brady (?).	× „ <i>micheliniana</i> (d'Orbigny).
×	<i>Globigerina æquilateralis</i> , Brady.	„ <i>partschiana</i> (d'Orbigny).
×	„ <i>bulloides</i> , d'Orbigny.	<i>Rotalia orbicularis</i> , d'Orbigny.
×	„ <i>conglobata</i> , Brady.	<i>Nonionina umbilicatula</i> (Montagu).
×	„ <i>dubia</i> , Egger.	

Cælotholus cruciatus was observed among the Radiolaria in the deposit from 1200 fathoms (Station 164A).

ORGANISMS FROM
SURFACE-NETS.

Surface Organisms.—The following two species of Radiolaria are recorded from the surface :—

RADIOLARIA (Hæeckel, Zool. pt. 40).

Dictyocephalus australis, Hæeckel.*Castanidium bromleyi*, Hæeckel.

In addition, the following are recorded in the note-books :—Foraminifera (a few specimens of *Orbulina* and of *Globigerina*), *Sagitta*, and many small Crustacea.