

*Caryophyllia lamellifera*, n.sp. Two specimens (630 fathoms); obtained at no other locality. STATIONS 170 AND 170A.

## CRINOIDEA (Carpenter, Zool. pts. 32 and 60).

*Pentacrinus naresianus*, n.sp. Two specimens and fragment (520 and 630 fathoms); obtained also at Stations 171, 175, 210 (?), and 214, 375 to 1350 fathoms.

*Metacrinus varians*, n.g., n.sp. (?). One specimen (630 fathoms); obtained also at Station 214, 500 fathoms.

„ *wyvillii*, n.g., n.sp. Two specimens (630 fathoms); obtained also at Station 214, 500 fathoms.

„ *nodosus*, n.g., n.sp. Three specimens (630 fathoms); obtained at no other locality.

*Antedon incerta*, n.sp. One specimen (630 fathoms); obtained at no other locality.

„ *breviradia*, n.sp. Five specimens (630 fathoms); obtained also at Station 175, 1350 fathoms.

„ *echinata*, n.sp. One specimen (630 fathoms); obtained at no other locality.

„ *basicurva*, n.sp. Several specimens (630 fathoms); obtained also at Station 175 (?), 1350 fathoms.

„ *incisa*, n.sp. Four specimens and fragments (630 fathoms); obtained also at Station 174, 210, 255, or 610 fathoms.

„ *alternata*, n.sp. Two specimens (630 fathoms); obtained also at Stations 169, 218, and 236.

„ *inæqualis*, n.sp. Twelve specimens (630 fathoms); obtained also at Stations 174 and 175, 210 to 1350 fathoms.

## ASTEROIDEA (Sladen, Zool. pt. 51).

*Solaster torulatus*, n.sp. (520 fathoms); obtained at no other locality.

*Cribrella sufflata*, n.sp. One specimen (520 fathoms); obtained at no other locality.

## OPHIUROIDEA (Lyman, Zool. pt. 14).

*Ophioceramis* (?) *clausa*, n.sp. (630 fathoms); obtained also at Station 171, 600 fathoms.

„ (?) *obstricta*, n.sp. (630 fathoms); obtained also at Station 192, 129 fathoms.

*Ophiactis cuspidata*, n.sp. (520 fathoms); obtained also at Station 171, 600 fathoms.

*Ophiacantha cornuta*, n.sp. (520 fathoms); obtained also at Station 171, 600 fathoms.

*Ophiomitra plicata*, n.sp. (630 fathoms); obtained also at Stations 205 and 214, 1050 and 500 fathoms.