

- Actinometra typica* (Lovén). One specimen (255, 610, or 210 fathoms); obtained at no other locality by the Challenger. Recorded from Malacca, Jobie, Zebu, Fiji, and Kingsmill Islands. STATIONS 174 TO 174D.
- „ *stelligera*, n.sp. Seven specimens (255, 610, or 210 fathoms—most probably 210 or 255 fathoms); obtained at no other locality by the Challenger. Recorded from Tonga, Fiji, Samoa, and Atagar.
- „ *parvicirra* (Müller). One specimen (255, 610, or 210 fathoms); for distribution, see the Cape.

OPHIUROIDEA (Lyman, Zool. pt. 14).

- Ophiophyllum petilum*, n.g., n.sp. (210 to 610 fathoms); obtained also at Station 171. Only species of the genus.
- Ophiactis nama*, n.sp. (210 to 610 fathoms); obtained also at Station 171.
- Amphiura bellis*, n.sp. (var. ?). (210 to 610 fathoms); obtained also at Stations 232 and 236, 345 to 775 fathoms.

ECHINOIDEA (Agassiz, Zool. pt. 9).

- Micropyga tuberculata*, n.g., n.sp. (255, 610, or 210 fathoms); obtained also at Stations 204, 219, and Philippines, 100 to 150 fathoms. Only species of the genus.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

- Synapta challengerii*, n.sp. Two specimens (210 fathoms); obtained at no other locality. [Recorded in error from 140 fathoms].

ANNELIDA (M'Intosh, Zool. pt. 34).

- Praxilla köllikeri*, n.sp. One specimen (210 fathoms); obtained at no other locality. [Recorded in error from 140 fathoms].
- Terebellides ehlersi*, n.sp. One specimen (210 fathoms); obtained at no other locality.

MYZOSTOMIDA (Graff, Zool. pts. 27 and 61).

- Myzostoma labiatum*, n.sp. (On *Antedon inæqualis* ?).
- „ *echinus*, n.sp. (On *Antedon incisa* or *A. inæqualis*).
- „ *elongatum*, Graff. (On *Antedon occulta*, *A. similis*, or *A. tuberculata*).
- „ *fissum*, n.sp. (On *Antedon inæqualis*, *A. occulta*, *A. similis*, or *A. tuberculata*).