CAPE YORK.

- Craniella simillima (Bowerbank), var. One specimen (Station 186, 8 fathoms); obtained also at Port Jackson and Philippines.
- Myriastra simplicifurca, n.g., n.sp. One specimen (Station 186, 8 fathoms); obtained at no other locality.
 - clavosa (Ridley). Several specimens (Station 186, 8 fathoms); obtained also at Stations 208 and 212, 8 and 10 fathoms. Recorded from Torres Strait and Arafura Sea.
- Pilochrota purpurea (Ridley), var. longancora, nov. One specimen (Torres Strait, 3 to 11 fathoms); obtained at no other locality by the Challenger. Recorded from Torres Strait and Arafura Sea.
 - , moseleyi, n.g., n.sp. One specimen (Torres Strait, 3 to 11 fathoms); obtained at no other locality.
- Disyringa dissimilis (Ridley). (Torres Strait, 3 to 11 fathoms); obtained also at Station 188, 28 fathoms. Recorded from Port Darwin and Torres Strait.
- Tethya seychellensis (Wright). One specimen (Station 186, 8 fathoms); obtained also at Philippines. Recorded from Seychelle Islands.

CALCAREA (Poléjaeff, Zool. pt. 24).

Leucosolenia challengeri, n.sp. One specimen (Station 186, 8 fathoms); obtained at no other locality.

Heteropegma nodus gordii, n.g., n.sp. One specimen (Station 186, 8 fathoms); obtained also at Station 36. Only species of the genus.

Anamixilla torresi, n.g., n.sp. One specimen (Torres Strait, 3 to 11 fathoms); obtained at no other locality. Only species of the genus.

Leuconia dura, n.sp. One specimen (Station 186, 8 fathoms); obtained also at Station 36.

ALCYONARIA (Wright and Studer, Zool. pts. 64 and 81).

- Echinogorgia pseudosassapo, Kölliker. One specimen (Torres Strait); obtained at no other locality by the Challenger. Recorded from Indo-Pacific region.
- Plexauroides prælonga (Ridley). Two specimens (Stations 186, 8 fathoms); obtained at no other locality by the Challenger. Recorded from Torres Strait.
- Leptogorgia torresia, n.sp. One specimen (Torres Strait); obtained at no other locality.