

- Ophiothrix stelligera*, Lyman. (Station 186, 8 fathoms); obtained also at Fiji, CAPE YORK.
Arafura Sea (?), and Philippines.
- Ophiomaza cacaotica*, Lyman. (Cape York, and Station 187, 6 fathoms); obtained
at no other locality by the Challenger.
- Euryale aspera*, Lamarck. (Station 186, 8 fathoms); obtained also at Station 203,
12 to 20 fathoms.

ECHINOIDEA (Agassiz, Zool. pt. 9).

- Phyllacanthus annulifera* (Lamarck). (Cape York, and Station 186, 8 fathoms);
obtained also at Station 188, 28 fathoms.
- „ *verticillata* (Lamarck). (Station 186, 8 fathoms); obtained at no
other locality by the Challenger.
- Pleurechinus bothyroides*, Agassiz. (Station 186, 8 fathoms); obtained also
at Japan.
- Salmacis globator*, Agassiz. (Station 186, 8 fathoms); obtained at no other
locality by the Challenger.
- „ *rarispinga*, Agassiz. (Station 186, 8 fathoms); obtained also at
Stations 188 and 203, 12 to 28 fathoms.
- Laganum depressum*, Lesson. (Station 187, 6 fathoms); obtained also in Arafura Sea.
- Peronella decagonalis* (Lesson). (Station 186, 8 fathoms, and Torres Strait); for
distribution see Station 172.
- Breynia australasiæ* (Leach). (Torres Strait); obtained at no other locality by
the Challenger.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

- Synapta verrilli*, n.sp. Two specimens (Station 186, 8 fathoms); obtained at no
other locality.
- Colochirus challengeri*, n.sp. Three specimens (Station 186, 8 fathoms); obtained
at no other locality.
- Actinocucumis typica*, Ludwig. One specimen (Station 186, 8 fathoms); obtained
at no other locality by the Challenger. Recorded from
Australia and China.
- Thyone fusus* (Müller), var. *papuensis*, nov. One specimen (Station 186, 8 fathoms);
obtained at no other locality. The species recorded from
European seas.

ENTOZOA (Linstow, Zool. pt. 71).

- Tænia clavulus*, n.sp. Numerous fragments (from intestine of *Ptilorhis alberti*,
Cape York).