

*Cælopleurus maillardi* (Michelin). Obtained also at Station 201 and Amboina, STATION 192.  
82 to 102 fathoms.

*Asthenosoma pellucidum*, n.sp. Obtained also at Station 209, 100 fathoms.

*Temnopleurus hardwicki* (Gray). Obtained also at Japan and Arafura Sea.

„ *reynaudi*, Agassiz. Obtained also at Stations 166 and 219.

*Peronella decagonalis* (Lesson). For distribution see Station 172.

*Echinolampas oviformis* (Gmelin). One specimen; obtained at no other locality  
by the Challenger.

*Catopygus recens*, n.sp. Obtained at no other locality.

*Maretia alta*, Agassiz. Obtained also at Station 191.

*Moiropsis claudicans*, n.g., n.sp. Obtained at no other locality. Only species of  
the genus.

#### HOLOTHURIOIDEA (Théel, Zool. pts. 13 and 39).

*Ilyodæmon maculatus*, n.g., n.sp. Two specimens; obtained also at Stations 209  
and 219, 95 to 150 fathoms. Only species of the genus.

*Chirodota* sp. (?). Several fragmentary specimens.

#### ANNELIDA (M'Intosh, Zool. pt. 34).

*Syllis ramosa*, n.sp. Obtained also at Station 209, 95 fathoms.

#### MYZOSTOMIDA (Graff, Zool. pts. 27 and 61).

*Myzostoma wyville-thomsoni*, n.sp. (On *Metacrinus angulatus*).

„ *folium*, n.sp. (On *Antedon manca*).

„ *inflator*, n.sp. (On *Antedon angustiradia*).

„ *murrayi*, n.sp. (On *Antedon angustiradia*).

„ *willemoesii*, n.sp. (On *Antedon flexilis*).

„ *beardi*, n.sp. (On *Antedon flexilis*).

#### CIRRIPIEDIA (Hoek, Zool. pt. 25).

*Lepas anserifera*, Linné. Several specimens (on branch of tree); widely distributed  
in Atlantic and Pacific Oceans.

*Scalpellum indicum*, n.sp. One specimen; obtained at no other locality.

„ *balanoides*, n.sp. Several specimens; obtained at no other locality.

#### MACRURA (Spence Bate, Zool. pt. 52).

*Ibaccus brevipes*, n.sp. One specimen; obtained at no other locality.

*Arctus sordidus*, Stimpson. Three specimens; obtained at no other locality by the  
Challenger. Recorded from Hong Kong.