

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Synapta beselii, Jæger. Several specimens; obtained also at Station 177, Philippines, and Tahiti.

Holothuria atra, Jæger. One specimen; obtained also at Station 177 and Fiji.

ANNELIDA (M'Intosh, Zool. pt. 34).

Nothria willemoesii, n.sp. One specimen (100 fathoms); obtained at no other locality.

Glycera amboinensis, n.sp. One specimen (15 fathoms); obtained at no other locality.

OSTRACODA (Brady, Zool. pt. 3).

Macrocypis maculata, Brady. (15 to 20 fathoms); obtained also at Stations 145, 149, 162, and Simon's Bay, Cape.

Bairdia amygdaloides, Brady. (15 to 20 fathoms); for distribution see Station 162.

Cythere scutigera, Brady. (15 to 20 fathoms); obtained also at Stations 167, 300, and New Guinea.

Cytheridea spinulosa, Brady. (15 to 20 fathoms); obtained also at Station 279, 420 fathoms. Recorded from Mauritius.

Cytherella pulchra, Brady. (15 to 20 fathoms); obtained also at Stations 33, 167, 344, and Port Jackson.

STOMATOPODA (Brooks, Zool. pt. 45).

Squilla chlorida, n.sp. One specimen (15 fathoms); obtained at no other locality.

Lysiosquilla maculata (Fabricius). Two specimens; obtained also at Philippines.

MACRURA (Spence Bate, Zool. pt. 52).

Scallasis amboinæ, n.g., n.sp. One specimen; obtained at no other locality. Only species of the genus.

Nika processa, n.sp. One specimen (15 fathoms); obtained at no other locality.

ANOMURA (Henderson, Zool. pt. 69).

Raninoides personatus, White, MS. Several specimens (15 fathoms); obtained at no other locality by the Challenger. Recorded from Eastern seas.

Cosmonotus grayii, Adams and White. One specimen (15 fathoms); obtained at no other locality by the Challenger. Recorded from Borneo and Formosa.

Munida spinulifera, Miers. One specimen (15 fathoms); obtained at no other locality by the Challenger. Recorded from Arafura Sea.

„ *militaris*, n.sp. One specimen (100 fathoms); obtained also at Stations 173, 192, 200 (var.), and 210 (var.).