

STATIONS 204
TO 204B.

trawl, which came up at 5 P.M. with numerous specimens. Proceeded through the channel between Simara and Banton Islands. Lightning with passing showers at night.

ANIMALS FROM
TRAWL.

The following species are recorded in the Zoological Reports from the trawlings in 100 to 115 fathoms (Stations 204A and 204B):—

ASTEROIDEA (Sladen, Zool. pt. 51).

Astropecten imbellis, n.sp. Obtained at no other locality.

Luidia aspera, n.sp. Obtained also at Station 219 and Samboangan, 150 and 10 fathoms.

Nymphaster symbolicus, n.g., n.sp. Obtained also at Stations 188 and 192 (var.).

Leptogonaster cristatus, n.g., n.sp. Obtained at no other locality. Only species of the genus.

Pholidaster squamatus, n.g., n.sp. Obtained at no other locality. The only other species of the genus (*Pholidaster distinctus*) was taken at Station 192.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiocamax vitrea, n.g., n.sp. Obtained also at Stations 192, 201, 209, and 219.

Ophiothrix capillaris, n.sp. Obtained also at Station 209, 100 fathoms.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Dorocidaris (Cidaris) papillata, Leske. For distribution see Station 24.

Goniocidaris florigera, n.sp. Obtained also at Station 192.

Micropyga tuberculata, n.g., n.sp. Obtained also at Stations 174, 209, and 219.
Only species of the genus.

Asthenosoma tessellatum, n.sp. One specimen; obtained at no other locality.

CIRRIPEDIA (Hoek, Zool. pt. 25).

Megalasma striatum, n.g., n.sp. Eight specimens (attached to Echini spines); obtained at no other locality. Only species of the genus.

Scalpellum rubrum, n.sp. One specimen; obtained at no other locality.

Balanus tenuis, n.sp. Four specimens; obtained at no other locality.

STOMATOPODA (Brooks, Zool. pt. 45).

Squilla leptosquilla, n.sp. One specimen; obtained at no other locality.

MACRURA (Spence Bate, Zool. pt. 52).

Nephrops thomsoni, n.sp. One specimen; obtained also at Station 166.

Penæus fissurus, n.sp. Three specimens; obtained also at Stations 190 and 209.