

STATION 209.

gigantic *Stylifer*. Only one Lepidid Cirriped was found, attached to *Molpadia*. There were beautiful Holothurians with wheels (large and small) like those of *Chirodota* distributed in whitish patches all over the body. Very fine specimens of *Sabellida* came up in long mud tubes. The Crustacea were represented by a particularly fine Stenorhynchid with long spiny legs, most of which were, however, broken off, and two other forms with one pair of legs on their backs, allied to the genus *Cymopolia*."

ORGANISMS FROM
THE DEPOSIT.

DIATOMACEÆ.—The following species of Diatoms were observed by Mr. Comber in the deposit from this Station (see also Murray and Renard, Deep-Sea Deposits Chall. Exp.) :—

<i>Navicula lyra</i> , Ehrenberg.	<i>Asteromphalus elegans</i> , Greville (= <i>A. wyvillei</i> , Castracane).
„ <i>clavata</i> , Gregory.	„ <i>roperianus</i> , Ralfs.
„ <i>prætexta</i> , Ehrenberg.	<i>Stictodiscus affinis</i> , Castracane.
„ <i>bullata</i> , Norman.	„ <i>californicus</i> , Greville.
„ <i>brasiliensis</i> , Grunow.	<i>Cyclotella stylorum</i> , Brightwell.
„ <i>indica</i> , Greville.	<i>Coscinodiscus excentricus</i> , Ehrenberg.
„ <i>baileyana</i> , Grunow.	„ <i>leptopus</i> , Grunow.
„ <i>durandii</i> , Kitton.	„ <i>nitidulus</i> , Gregory.
„ <i>nitescens</i> , Ralfs.	„ <i>nitidulus</i> , Grunow.
„ <i>bombus</i> , Kutzing.	„ <i>marginatus</i> , Ehrenberg.
„ <i>pandura</i> , Brebisson.	„ <i>subtilis</i> , Ehrenberg.
„ <i>aspera</i> , Ehrenberg.	„ <i>curvatus</i> , Grunow.
„ sp. (?).	„ var. <i>subocellatus</i> , Grunow.
<i>Alloioneis antillarum</i> , Cleve and Grunow.	„ <i>radiatus</i> , Ehrenberg.
<i>Pleurosigma affine</i> , Grunow, var. <i>normanni</i> , Ralfs.	„ <i>apolinisi</i> , Ehrenberg.
„ <i>naviculacea</i> , Brebisson.	„ <i>africanus</i> , Janisch.
<i>Nitzschia vermiculata</i> , Castracane.	„ „ var. <i>wallichiana</i> , Grunow.
„ <i>obesa</i> , Castracane.	„ <i>apiculatus</i> , Ehrenberg.
„ <i>panuliformis</i> , Gregory.	„ <i>nodulifer</i> , Janisch.
„ <i>plana</i> , Smith.	„ <i>reniformis</i> , Castracane (= <i>Stoschia admirabilis</i> , Janisch).
<i>Synedra robusta</i> , Ralfs.	<i>Pyxidicula cruciata</i> , Ehrenberg.
<i>Surirella fastuosa</i> , Ehrenberg.	<i>Pyxilla</i> sp. (?).
„ <i>recedens</i> , A. Schmidt.	<i>Paralia sulcata</i> , Cleve.
„ <i>dives</i> , Castracane.	<i>Stephanopyxis superba</i> , Grunow.
<i>Campylodiscus intermedius</i> , Greville.	<i>Podosira ambigua</i> , Grunow.
„ <i>robertsonianus</i> , Greville.	„ <i>argus</i> , Grunow.
„ <i>ambiguus</i> , Greville (= <i>C. latus</i> , Shadbolt).	<i>Hemidiscus cuneiformis</i> , Wallich.
„ <i>biangulatus</i> , Greville.	<i>Triceratium robertsonianum</i> , Greville.
„ <i>ornatus</i> , Greville.	„ <i>latum</i> , Greville.
„ <i>kittonianus</i> , Greville.	„ <i>favus</i> , Ehrenberg.
„ <i>undulatus</i> , Greville.	„ <i>jimbriatus</i> , Ehrenberg.
„ <i>angulatus</i> , Gregory.	„ sp. (?).
<i>Actinocyclus pruinosus</i> , Castracane.	<i>Biddulphia tuomeyii</i> , Roper.
„ <i>clevei</i> , Castracane.	„ <i>reticulata</i> , Roper.
„ sp. (?).	„ sp. (?).
<i>Actinoptychus areolatus</i> , A. Schmidt.	
<i>Asteromphalus brookei</i> , Bailey.	