

## PHILIPPINES.

- Ophiactis savignyi* (Müller and Troschel). (Samboangan, 10 fathoms); obtained also at Station 208.
- Ophiocoma scolopendrina* (Lamarck). (Samboangan Bank and Zebu reefs); obtained also at the Cape, Tongatabu, and Fiji.
- „ *erinaceus*, Müller and Troschel. (Samboangan Bank); obtained also at Sandwich Islands.
- Ophiomastix caryophyllata*, Lütken. (Zebu reefs); obtained also at Fiji.
- Ophiothrix longipeda* (Lamarck). (Samboangan, 10 fathoms); for distribution see Station 172.
- „ *ciliaris* (Lamarck) (?). (Zebu); obtained also at Station 209.
- „ *stelligera*, Lyman. (Samboangan); obtained also at Station 186, Fiji, and Arafura Sea.
- „ *propinqua*, Lyman (?). (Samboangan); obtained also at Station 172 (?) and Fiji.
- „ *nereidina* (Lamarck). (Samboangan, 10 fathoms); obtained at no other locality by the Challenger.
- „ *striolata*, Grube. (Samboangan, 10 fathoms); obtained also at Station 208.

## ECHINOIDEA (Agassiz, Zool. pt. 9).

- Phyllacanthus baculosa* (Lamarck). (Samboangan, 10 fathoms); obtained also at station 201.
- Diadema setosum*, Gray. (Zebu); obtained also at Cape Verdes, St. Thomas, and Tahiti.
- Echinothrix calamaris* (Pallas). (Zebu reefs); obtained also at Fiji and Tahiti.
- Asthenosoma grubii*, n.sp. (Samboangan, 10 fathoms); obtained at no other locality.
- Echinometra lucunter* (Leske). (Samboangan and Zebu, 10 fathoms); for distribution see Station 172.
- Pseudoboletia indiana* (Michelin). (Samboangan, 10 fathoms); obtained at no other locality by the Challenger.
- Salmacis bicolor*, Agassiz. (Samboangan, 10 fathoms); obtained at no other locality by the Challenger.
- „ *sulcata*, Agassiz. (Zebu); obtained at no other locality by the Challenger.
- Mespilia globulus*, Agassiz. (Samboangan, 10 fathoms); obtained at no other locality by the Challenger.
- Toxopneustes pileolus* (Lamarck). (Samboangan, 10 fathoms); obtained also at Sandwich Islands.
- Hipponoë variegata* (Leske). (Samboangan, 10 fathoms); obtained also at Port Jackson and Fiji.