

ADMIRALTY  
ISLANDS.

- Cythere obtusalata*, n.sp. (16 to 25 fathoms); obtained also at Station 162.  
 „ *prava* (Baird). (16 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Mediterranean.  
 „ *irrorata*, n.sp. (16 to 25 fathoms); obtained at no other locality.  
 „ *tricristata*, n.sp. (16 to 25 fathoms); obtained also at Port Jackson.  
*Loxococoncha pumicosa*, n.sp. (16 fathoms); obtained also at Station 187.  
*Xestoleberis tumefacta*, n.sp. (16 fathoms); obtained at no other locality.  
*Cytherella semitalis*, Brady. (16 fathoms); for distribution see Station 187.

## ANOMURA (Henderson, Zool. pt. 69).

- Cænobita clypeata* (Herbst). One specimen (Wild Island); obtained at no other locality by the Challenger. Recorded from Indo-Pacific region.  
 „ *rugosa*, M.-Edwards. Several specimens (Tracey and Wild Islands); obtained also at Fiji, New Hebrides, Arrou Islands, and Tahiti.  
*Pagurus dearmatus*, n.sp. One specimen (16 to 25 fathoms); obtained at no other locality.  
*Spiropagurus spiriger* (De Haan). One specimen (16 to 25 fathoms); for distribution see Torres Strait.

## BRACHYURA (Miers, Zool. pt. 49).

- Neptunus (Achelous) granulatus* (M.-Edwards). One specimen (16 to 25 fathoms); obtained also at Philippines.  
*Cardiosoma carnifex* (Herbst). Four specimens (Tracey Island); obtained also at Fiji and Tahiti.  
*Sesarma (Holometopus) aubryi*, M.-Edwards. Several specimens (Wild Island, beach); obtained also at New Hebrides and Arrou Islands.

## LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

- Mactra achatina*, Chemnitz. (16 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Indian and Pacific Oceans.  
 „ *incerta*, n.sp. (16 to 25 fathoms); obtained at no other locality.

## GASTEROPODA (Watson, Zool. pt. 42).

- Trochus (Chrysostoma) paradoxus* (Born). (Wild Island); obtained also at Philippines.  
*Vitulularia crenifera* (Montrouzier). (16 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from New Caledonia.  
*Harpa nobilis*, Lamarek. (16 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Pacific islands.  
*Cassis (Casmaria) turgida*, Reeve. (16 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Philippines.