

STATION 225.

II. NASSELLARIA.—Of the Nassellaria, the Plectoidea, Stephoidea, and Botryodea are but sparsely represented, while the Spyroidea and Cyrtodea are abundant and large; in fact, the individuals belonging to the Spyroidea and Cyrtodea make up three-fourths of the Nassellaria, and one-half of the total Radiolaria, present in the deposit.

- | | |
|---|--|
| <i>Clathrocircus multiformis</i> , Haeckel. | <i>Phormospyris tridentata</i> , Haeckel. |
| <i>Zygostephanus reniformis</i> , Haeckel. | <i>Desmospyris cyrtocolpos</i> , Haeckel. |
| <i>Coronidium dyostephanus</i> , Haeckel. | " <i>mammillata</i> , Haeckel. |
| <i>Triplagia horizontalis</i> , Haeckel. | <i>Tricolospyris kantiana</i> , Haeckel. |
| <i>Archicircus ellipsis</i> , Haeckel. | " <i>leibnitziana</i> , Haeckel. |
| <i>Lithocircus crambessa</i> , Haeckel. | " <i>newtoniana</i> , Haeckel. |
| <i>Dendrocircus stalactites</i> , Haeckel. | <i>Nephrospyris coriata</i> , Haeckel. |
| <i>Semantis dipyla</i> , Haeckel. | <i>Perispyris bicincta</i> , Haeckel. |
| <i>Semantiscus hexapylus</i> , Haeckel. | <i>Botryopera multiloba</i> , Haeckel. |
| <i>Eucoronis laevigata</i> , Haeckel. | <i>Cannobotrys pentacanna</i> , Haeckel. |
| <i>Acrocubus octopylus</i> , Haeckel. | <i>Botryopyle cephalodes</i> , Haeckel. |
| " <i>tesseralis</i> , Haeckel. | " <i>sethocorys</i> , Haeckel. |
| <i>Toxarium bifurcum</i> , Haeckel. | <i>Botryocampe camerata</i> , Haeckel. |
| <i>Ocotympanum arborescens</i> , Haeckel. | " <i>rotalia</i> , Haeckel. |
| <i>Prismatium tripodium</i> , Haeckel. | <i>Lithobotrys sphærothorax</i> , Haeckel. |
| <i>Tympanidium binoctonum</i> , Haeckel. | <i>Pylobotrys fontinalis</i> , Haeckel. |
| <i>Lithotympanum tuberosum</i> , Haeckel. | <i>Phormobotrys cannothalamia</i> , Haeckel. |
| <i>Eutympanium militare</i> , Haeckel. | " <i>pentathalamia</i> , Haeckel. |
| <i>Tripospyris conifera</i> , Haeckel. | <i>Trissopilium lithomelissa</i> , Haeckel. |
| " <i>diomma</i> , Haeckel. | <i>Tripilidium ovatum</i> , Haeckel. |
| " <i>semantis</i> , Haeckel. | <i>Tripodiscium tristyllospyris</i> , Haeckel. |
| " <i>tribrachiata</i> , Haeckel. | <i>Euscenium archicyrtes</i> , Haeckel. |
| <i>Triceruspyris corallorrhiza</i> , Haeckel. | <i>Cladioscenium gladiatum</i> , Haeckel. |
| " <i>giraffa</i> , Haeckel. | " <i>pinnatum</i> , Haeckel. |
| <i>Tristyllospyris scaphipes</i> , Haeckel. | <i>Pteroscenium spinulosum</i> , Haeckel. |
| " <i>ramosa</i> , Haeckel. | <i>Peridium cervinum</i> , Haeckel. |
| <i>Dipospyris cubus</i> , Haeckel. | <i>Bothropyramis quadrata</i> , Haeckel. |
| " <i>forcipata</i> , Haeckel. | <i>Carpocanistrum giganteum</i> , Haeckel. |
| " <i>sigmopodium</i> , Haeckel. | <i>Cornutella mitra</i> , Haeckel. |
| <i>Dendrospyris furcata</i> , Haeckel. | <i>Archiscenium callimitra</i> , Haeckel. |
| <i>Dorcadospyris antilope</i> , Haeckel. | <i>Archipera cortiniscus</i> , Haeckel. |
| <i>Elaphrosphyris capricornis</i> , Haeckel. | <i>Cyrtophormis spiralis</i> , Haeckel. |
| " <i>cervicornis</i> , Haeckel. | <i>Cornutanna loxoconus</i> , Haeckel. |
| <i>Petalospyris eupetala</i> , Haeckel. | <i>Archicorys ampulla</i> , Haeckel. |
| <i>Liriospyris heteropoda</i> , Haeckel. | " <i>microstoma</i> , Haeckel. |
| <i>Therospyris leo</i> , Haeckel. | <i>Halicapsa hystrix</i> , Haeckel. |
| <i>Ceratospyris allmersii</i> , Haeckel. | <i>Cyrtocalpis gromia</i> , Haeckel. |
| " <i>carnerii</i> , Haeckel. | <i>Dictyophimus bicornis</i> , Haeckel. |
| " <i>strasburgeri</i> , Haeckel. | " <i>brandtii</i> , Haeckel. |
| <i>Gorgospyris schizopodia</i> , Haeckel. | " <i>challengeri</i> , Haeckel. |
| <i>Dictyospyris mammillaris</i> , Haeckel. | " <i>sphærocephalus</i> , Haeckel. |
| <i>Lophospyris dipodiscus</i> , Haeckel. | " <i>triserratus</i> , Haeckel. |
| <i>Tiarospyris mitra</i> , Haeckel. | <i>Sethopilium orthopus</i> , Haeckel. |
| " <i>pervia</i> , Haeckel. | <i>Psilomelissa calvata</i> , Haeckel. |