

<i>Coscinodiscus curvatus</i> , Grunow.
" " var. <i>recta</i> , Rattray.
" <i>radiatus</i> , Ehrenberg.
" " var. <i>media</i> , Grunow.
" <i>nodulifer</i> , A. Schmidt.
" <i>obscurus</i> , A. Schmidt.
" <i>africanus</i> , Janisch.
" <i>concinus</i> , Smith.
" <i>rex</i> , Wallich.
" <i>centralis</i> , Ehrenberg.
" <i>marginatus</i> , Ehrenberg.
" <i>normanii</i> , Gregory.
" <i>polyacanthus</i> , Grunow (?).
<i>Coscinodiscus dubiosus</i> , Grunow.
" <i>gazellæ</i> , Janisch.
<i>Ethmodiscus</i> sp. (?).
<i>Actinocyclus punctulatus</i> , Castracane.
" sp. (near <i>subtilis</i>).
<i>Asteromphalus arachne</i> , Ralfs.
" <i>roperianus</i> , Ralfs.

<i>Asteromphalus humboldtii</i> , Ehrenberg.
" <i>hiltonianus</i> , Ralfs.
<i>Stoschia admirabilis</i> , Janisch.
<i>Asterolampra marylandica</i> , Ehrenberg.
<i>Melosira</i> sp. (?).
<i>Surirella robusta</i> , Ehrenberg (fresh water), var.
<i>Fragilaria pacifica</i> , Grunow.
" <i>nitzschiooides</i> , Grunow, var. <i>brasiliensis</i> , Rattray, MS.
<i>Achnanthes subsessilis</i> , Ehrenberg.
<i>Cymbella (Cocconema) cistula</i> , Hempricht (fresh water).
<i>Synedra capitata</i> , Ehrenberg (fresh water).
" <i>gaillionii</i> , Ehrenberg.
<i>Epithemia turgida</i> , Kutzing, var. <i>granulata</i> , Cleve (fresh water).
" " forma minor.
" <i>gibba</i> , Kutzing, var. <i>ventricosa</i> , Cleve (fresh water).
<i>Pleurosigma decorum</i> , W. Smith.
<i>Gomphonema geminatum</i> , Agardh (fresh water).

STATION 229.

Surface Organisms.—The following species are recorded from the surface in the neighbourhood of this Station :—

ORGANISMS FROM SURFACE-NETS.

RADIOLARIA (Haeckel, Zool. pt. 40).

- Sethoconus trochus*, Haeckel.
Theocampe cryptoprora, Haeckel.

SIPHONOPHORÆ (Haeckel, Zool. pt. 77).

- Velella oblonga*, Chamisso.

AMPHIPODA (Stebbing, Zool. pt. 67).

- Synopia schéeleana*, Bovallius.
Vibilia propinquæ, n.sp.
Phrosina pacifica, n.sp.
Anchylomera sp. (?).

Phorcorrhaphis edwardsi, n.sp.*Platyscelus armatus* (Claus).*Eupronoë* sp. (?).*Brachyscelus crusculum*, Spence Bate.

CEPHALOPODA (Hoyle, Zool. pt. 44).

- Octopus brevipes*, d'Orbigny.
Teleoteuthis caribbæa (Lesueur), Verrill (?).

TUNICATA (Herdman, Zool. pt. 76).

- Doliolum affine*, n.sp.

In addition, the following are recorded in the note-books (April 1 to 4):—
Oscillatoriaceæ, *Peridinium*, *Pyrocystis*, *Ethmodiscus* and other Diatoms (the smaller species of Diatoms more abundant than in any of the recent Stations to the south), *Globigerina*, *Orbulina*, *Hastigerina*, *Pulvinulina*, *Calcaromma calcarea* and other Radiolaria in great numbers, Tintinnidæ and other Infusoria, Medusæ, *Diphyes*, *Porpita*, Ctenophoræ (*Cydippe* or *Pleurobrachia*), *Sagitta*, *Alciopa*, *Tomopteris*, Sipunculid larvæ, *Cypridina*, Copepods, *Oxycephalus*, *Phronima*, *Phronimella*, Hyperids, *Dithyrus*, *Erichthus*, *Alima*, *Euphausia pellucida*, *Siriella*, *Amphion*, *Phyllosoma*, *Lucifer*,