

STATION 235.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophioglypha lapidaria, n.sp. Obtained at no other locality.

Ophiomusium lymani, Wyville Thomson. For distribution see Station 45.

Ophiopholis japonica, n.sp. Obtained also at Station 236, 420 to 775 fathoms.

Amphiura acacia, n.sp. Obtained at no other locality.

Ophiacantha scolopendrica, n.sp. [See *Bull. Mus. Comp. Zool.*, vol. x. p. 259, 1883].

Ophiomitra normani, n.sp. Obtained also at Stations 232 and 236.

Ophiomyces spathifer, n.sp. Obtained at no other locality.

Astroceras pergamena, n.g., n.sp. Obtained at no other locality. Only species of the genus.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Phormosoma hoplacantha, Wyville Thomson, n.sp. Two specimens; obtained also at Stations 164B and 300.

Echinus norvegicus, Düben and Koren. Obtained also at Stations 46, 47, 232, and 308.

HOLOTHURIOIDEA (Théel, Zool. pts. 13 and 39).

Lætmogone spongiosa, n.g., n.sp. One specimen; obtained at no other locality.

Pælopatides (?) *appendiculata*, n.g., n.sp. Three specimens; obtained at no other locality.

ANNELIDA (M'Intosh, Zool. pt. 34).

Lætmonice aphroditoides, n.sp. One specimen; obtained at no other locality.

Polynoëlla levisetosa, n.g., n.sp. One specimen; obtained at no other locality.
Only species of the genus.

Hyalinæcia tubicola, Müller. One specimen; for distribution see Station VIII.

CIRRIPEDIA (Hoek, Zool. pt. 25).

Scalpellum japonicum, n.sp. One specimen; obtained at no other locality.

ISOPODA.

One specimen, genus and species undetermined.

MACRURA (Spence Bate, Zool. pt. 52).

Benthesicymus altus, n.g., n.sp. Four specimens; for distribution see Station 133.

Gennadas parvus, n.g., n.sp. One specimen; for distribution see Station 45.

Acanthephyra sica, n.sp. Two specimens; for distribution see Station 40.

Nematocarcinus tenuipes, n.sp. Six specimens; obtained also at Stations 218 and 232.

Notostomus japonicus, n.sp. One specimen; obtained at no other locality.