

SANDWICH
ISLANDS.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Stichopus godeffroyi, Semper, var. One specimen (Sandwich Islands); obtained at no other locality by the Challenger. Recorded from Navigator, Friendly, Fiji, and Pelew Islands.

Holothuria fusco-rubra, n.sp. One specimen (Sandwich Islands); obtained at no other locality.

OSTRACODA (Brady, Zool. pt. 3).

Pontocypris faba (Reuss) (?). (Honolulu reefs, 40 fathoms); obtained also at Station 162.

Bairdia amygdaloides, Brady. (Honolulu reefs, 40 fathoms); for distribution see Station 162.

„ *crosskeiana*, Brady. (Honolulu reefs, 40 fathoms); obtained also at Tongatabu and Admiralty Islands.

„ *expansa*, n.sp. (Honolulu reefs, 40 fathoms); obtained at no other locality.

„ *attenuata*, n.sp. (Honolulu reefs, 40 fathoms); obtained also at Station 185.

Cythere flos-cardui, n.sp. (Honolulu reefs, 40 fathoms); obtained at no other locality.

„ *reussi*, Brady. (Honolulu reefs, 40 fathoms); obtained also at Stations 187 and 313.

„ *rastromarginata*, n.sp. (Honolulu reefs, 40 fathoms); obtained also at Stations 162 and 167.

„ *quadriaculeata*, n.sp. (Honolulu reefs, 40 fathoms); obtained also at Station 233B.

„ *convoluta*, Brady. (Honolulu reefs, 40 fathoms); obtained also at Station 172.

Loxoconcha honoluluensis, n.sp. (Honolulu reefs, 40 fathoms); obtained at no other locality.

„ *alata*, Brady. (Honolulu reefs, 40 fathoms); obtained at no other locality by the Challenger. Recorded from Mauritius.

„ *anomala*, n.sp. (Honolulu reefs, 40 fathoms); obtained at no other locality.

Xestoleberis curta, Brady. (Honolulu reefs, 40 fathoms); for distribution see Station 33.

Cytherella venusta, n.sp. (Honolulu reefs, 40 fathoms); obtained at no other locality.

STOMATOPODA (Brooks, Zool. pt. 45).

Squilla nepa, Latreille. One specimen (Honolulu reefs); obtained also at Japan.

Pseudosquilla ciliata, Miers. One specimen (Honolulu reefs); obtained also at St. Thomas.