FALKLAND ISLANDS. Echinus magellanicus, Phil. (Station 315, 5 to 12 fathoms); for distribution see Station 145.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Chirodota purpurea, Lesson. Numerous specimens (Station 316, 4 to 5 fathoms); obtained at no other locality by the Challenger. Recorded from Falklands.

" australiana, Stimpson. Two specimens (Station 315, 5 to 10 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.

Cucumaria crocea, Lesson. Numerous specimens (Stations 315 and 316, 4 to 12 fathoms); obtained also at Stations 313 and 314.

mendax, n.sp. Six specimens (Station 315, 5 to 12 fathoms); obtained at no other locality.

Annelida (M'Intosh, Zool. pt. 34).

Hermadion kerguelensis, n.sp. Several specimens (Station 315, 5 to 10 fathoms); obtained also at Station 149.

Nereis (Platynereis) eatoni, M'Intosh. One specimen (Station 315, 5 to 10 fathoms); obtained also at Stations 113, 145, and 149.

OSTRACODA (Brady, Zool. pt. 3).

Aglaia (?) meridionalis, n.sp. (Stanley Harbour, 6 fathoms); obtained at no other locality.

Cythere moseleyi, n.sp. (Stanley Harbour, 6 fathoms); obtained at no other locality.

" falklandi, n.sp. (Stanley Harbour, 6 fathoms); obtained at no other locality.

,, fulvotincta, n.sp. (Stanley Harbour, 6 fathoms); obtained at no other locality.

" impluta, n.sp. (Stanley Harbour, 6 fathoms); obtained also at Station 135.

Xestoleberis polita, Brady. (Stanley Harbour, 6 fathoms); obtained at no other locality by the Challenger. Recorded from Magellan Strait.

Cytherura clavata, n.sp. (Stanley Harbour, 6 fathoms); obtained at no other locality.

AMPHIPODA (Stebbing, Zool. pt. 67).

Halirages huxleyanus (Spence Bate). One specimen (Stanley Harbour, from kelp); obtained at no other locality by the Challenger. Recorded from Hermit Island.