

Pylorena armata, Haeckel.
Spongotorchus parma, Haeckel.
Dictyocoryne echinata, Haeckel.
Stauracantha quadrifurca, Haeckel.
Quadrilonche telostaura, Haeckel.
Amphibelone aciculata, Haeckel.
Hystrichaspis sulcata, Haeckel.
Acontaspis lanceolata, Haeckel.
Phatnaspis cuscinoidea, Haeckel.

Carpocanistrum novenum, Haeckel.
Sethophormis floscula, Haeckel.
Setoconus orthoceras, Haeckel.
Clathrocyclas latonæ, Haeckel.
Aulographis boricornis, Haeckel.
 " *cruciata*, Haeckel.
Sayena triangula, Haeckel.
Challengeria thomsoni, Murray.

STATION 325.

OSTRACODA (Brady, Zool. pt. 3).

Halocypris atlantica, Lubbock.
 " *imbricata*, n.sp.

COPEPODA (Brady, Zool. pt. 23).

Calanus tonsus, n.sp.
Rhincalanus gigas, n.sp.
Pleuromma abdominalis (Lubbock).
Heterochaeta spinifrons, Claus.
Leuckartia flavigaster, Claus.
Undina darwinii, Lubbock.
Euchæta prestandreae, Philippi.
 " (?) *philippii*, n.sp.
 " *australis*, n.sp.
 " *gigas*, n.sp.
 " *barbata*, n.sp.

Phyllopodus bidentatus, n.g., n.sp.

Setella gracilis, Dana.
Corycaeus pellucidus, Dana.
 " *obtusus*, Dana.

Oncæa obtusa (Dana).
Nogagus lunatus, Steenstrup and Lütken, (parasitic on *Carcharias*).

TUNICATA (Herdman, Zool. pt. 76).

Salpa democratica - mucronata,
 Forskål.

FISHES (Günther, Zool. pt. 78).

Carcharias glaucus, Linné.

In addition, the following are recorded in the note-books:—Oscillatoriaceæ, *Peridinium tripos*, Diatoms, *Globigerina*, *Hastigerina*, Medusæ, Siphonophoræ, *Beroë*, *Sagitta*, *Alciopa*, Annelid larvae, Hyperids, Mysids and larvae, larvae of *Sergestes* and *Lucifer*, small crab, larval Gasteropods, *Atlanta* and other Heteropods, *Cavolinia*, *Styliola*, *Limacina*, and other Pteropods (with eggs as at the previous Station), *Appendicularia*, Scopelid, two inches in length, with brilliant scales, flying fish half an inch in length, and other small fishes. Mr. Murray was away in a boat, and noticed on the surface only very small *Velellæ*; with the water-glass many small things could be seen, but on the whole the water was very clear and blue. The tow-nets at the weights and trawl contained many surface and deep-sea organisms, including Challengeridæ and other Radiolaria, *Sagitta*, *Cypridina*, red Copepods, and a large red Schizopod.