

VIGO BAY.
ORGANISMS FROM
THE DEPOSIT.

OSTRACODA (Brady, Zool. pt. 3).

Paracypris polita, Sars.
Cythere tenera, Brady.
 „ *stimpsoni*, Brady.
Loxoconcha guttata (Gorman).

Cytheropteron intermedium, Brady.
Paradoxostoma ensiforme, Brady.
Polycope orbicularis, Sars.

ORGANISMS FROM
SURFACE-NETS.

Surface Organisms.—The following species is recorded from the surface in Vigo Bay :—

AMPHIPODA (Stebbing, Zool. pt. 67).

Gammarus locusta (Linné).

The tow-nets yielded in addition :—Many *Noctiluca*, *Peridinium*, Diatoms, Medusæ, *Evadne*, Copepods, Cirriped larvæ, Zoëæ, larval Lamellibranchs and Gasteropods, *Appendicularia*, and larval fishes.

NORTH ATLANTIC.

The following species are recorded from deep water dredgings (over 1000 fathoms) in the North Atlantic, the exact locality not being noted :—

ANIMALS FROM
DREDGE AND
TRAWL.

LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

Lucina (Codakia) tigerina (Linné). Obtained at no other locality by the Challenger.
Montacuta cylindræa, n.sp. Obtained at no other locality.

GASTEROPODA (Watson, Zool. pt. 42).

Clathurella streptophora, n.sp. Obtained at no other locality.
Natica radiata, n.sp. Obtained also at Station 33.
Mitrularia uncinata (Reeve). Obtained also at Stations 113A and 122.
Siliquaria (Tenagodus) bernardii, Mörch. Obtained at no other locality by the Challenger. Recorded from Senegal.

ORGANISMS FROM
SURFACE NETS.

The following species are recorded from the surface of the North Atlantic, the exact locality not being noted :—

AMPHIPODA (Stebbing, Zool. pt. 67).

Amphithyrus bispinosus, Claus.

MACRURA (Spence Bate, Zool. pt. 52).

Mastigopus suhmi, n.sp.
Sergestes ovatoculus, n.sp.
 „ *penerinkii*, n.sp.

Lucifer typus, Thompson.
 „ *reynaudii*, M.-Edwards.
Hectarthropus tenuis, n.g., n.sp.

CEPHALOPODA (Hoyle, Zool. pt. 44).

Alloposus mollis, Verrill.