

Buskiella abyssorum.
Lætmonice producta.
Cythere acanthoderma.
 „ *dasyderma.*
 „ *dictyon.*
Cytheropteron mucronalatum.
Gnathophausia gigas.
Acanthephryra sica.
Aristeus armatus.
Benthescymus brasiliensis.
Gennadas parvus.
Hemipenaeus spinidorsalis.
Hymenodora mollicutis.
Pontophilus gracilis.
Willemæsia leptodactyla.
Parapagurus abyssorum.
Arca (Barbatia) pteroessa.

Eledone rotunda.
Bifuraria reticulata.
Bugula margaritifera.
 „ *reticulata.*
Farcimina delicatissima.
Discina atlantica.
Pyrosoma spinosum.
Bathysaurus mollis.
Gonostoma microdon.
Ipnops murrayi.
Macrurus armatus.
 „ *filicauda.*
 „ *liocephalus.*
Melanocetus murrayi.
Sternopyx diaphana.
Synaphobranchus bathybius.

The following 42 species (or 43 per cent.) were taken in the zone between 1000 and 1500 fathoms:—

Bathypterus patula.
Amphianthus ornatum.
Corallimorphus profundus.
 „ *rigidus.*
Bathyactis symmetrica.
Deltocyathus italicus.
Freyella echinata.
Ophiacantha cosmica.
Ophioglypha minuta.
Cystechinus wyrillii.
Benthodytes papillifera.
Cucumaria abyssorum.
Oneirophanta mutabilis.
Pælopatides confundens.
Pseudostichopus villosus.
Cythere acanthoderma.
 „ *dasyderma.*
 „ *dictyon.*
Cytheropteron mucronalatum.
Arcturus abyssicola.
Acanthephryra sica.

Aristeus armatus.
Benthescymus brasiliensis.
Gennadas parvus.
Petalidium foliaceum.
Pontophilus gracilis.
Willemæsia leptodactyla.
Parapagurus abyssorum.
Arca (Barbatia) corpulenta.
Dentalium acutissimum.
Cirroteuthis magna.
Enoplateuthis margaritifera.
Bugula reticulata.
Discina atlantica.
Terebratula wyrillii.
Corynascidia suhmi.
Octacnemus bythius.
Gonostoma microdon.
Macrurus armatus.
 „ *filicauda.*
Sternopyx diaphana.
Synaphobranchus bathybius.

The following 18 species (or 19 per cent.) were taken in the zone between 500 and 1000 fathoms:—

Bathyactis symmetrica.
Deltocyathus italicus.
Pontaster venustus.
Ophiacantha cosmica.

Cythere acanthoderma.
 „ *dasyderma.*
 „ *dictyon.*
Serolis neæra.