

FISHES:

* <i>Bathylagus antarcticus</i> , n.g., n.sp.	S
<i>Bathyonus compressus</i> , n.sp.	T
<i>Bathysaurus mollis</i> , n.g., n.sp.	T N
* <i>Chiasmodus niger</i> , Johnson.	T
* <i>Cyema atrum</i> , n.g., n.sp.	S
<i>Gonostoma microdon</i> , n.sp.	S T N
<i>Ipnops murrayi</i> , n.g., n.sp.	S T
* <i>Macrurus affinis</i> , n.sp.	S
,, <i>altipinnus</i> , n.sp.	N
,, <i>armatus</i> , Hector.	S T N
* ,, <i>asper</i> , n.sp.	N
,, <i>filicauda</i> , n.sp.	S

<i>Macrurus liocephalus</i> , n.sp.	N
<i>Melamphaës crassiceps</i> , n.sp.	S T
* ,, <i>robustus</i> , n.sp.	T
<i>Melanoctetus murrayi</i> , n.sp.	T
* <i>Melanonus gracilis</i> , n.g., n.sp.	S
* <i>Neobythites grandis</i> , n.sp.	N
* <i>Notacanthus rissoanus</i> , Filippi and Vérany.	N
* <i>Platypterus apus</i> , n.g., n.sp.	T
* <i>Scopelus antarcticus</i> , n.sp.	S
<i>Sternopyx diaphana</i> (Herm.).	S T N
* <i>Stomias boa</i> (Risso).	S
<i>Synaphobranchus bathybius</i> , n.sp.	S N

Notes on the Bathymetrical Distribution.—Of the species enumerated in the above list 61 species (or 15 per cent.) were taken at more than one Station (when taken at more than two Stations the number of times is indicated within brackets after the name), viz. :—

Thenea fenestrata.
Umbellula carpenteri.
Cereus spinosus.
Liponema multiporum.
Bathyactis symmetrica (4).
Leptopenus discus (3).
Hyocrinus bethellianus.
Promachocrinus abyssorum.
Lonchotaster forcipifer.
Pararchaster pedicifer.
Ophiacantha cosmica (5).
Ophioglypha loveni (3).
Ophiomusium armigerum.
Cystechinus clypeatus.
 „ *wyvillii* (3).
Goniocidaris canaliculata (3).
Pourtalesia carinata.
 „ *hispida*.
Salenia hastigera.
Spatagocystis challengerii.
Urechinus naresianus.
Benthodytes sordida (3).
Cucumaria abyssorum.
Elpidia purpurea.
Laetmogone wyville-thomsoni.
Psychropotes longicauda.

Pseudostichopus villosus (4).
Phascolion lutense.
Grubianella antarctica (3).
Lætmonice producta.
Polynoë (Robertianella) synophtalma.
Cythere dasyderma.
 „ *dictyon* (4).
Cytheropteron mucronatum (3).
Krithe producta.
Cystisoma spinosum.
Stenopleura atlantica.
Eurycope fragilis (3).
Benth euphausia amblyops.
Boreomysis scyphlops (3).
Encopia australis (4).
Acanthephyra acanthitelsonis.
 „ *purpurea*.
Aristeus armatus (3).
Glyphocrangon rimipes.
Hymenodora mollicutis (4).
Willemoesia leptodactyla.
Parapagurus abyssorum.
Phoxichilidium pilosum.
Arca (Barbatia) pteroessa.
Verticordia tornata.
Bicellaria infundibulata.