

<i>Boltenia legumen</i> (3).	<i>Lethrinus nebulosus.</i>
<i>Colella pedunculata</i> (5).	<i>Lichia glauca.</i>
<i>Corella japonica</i> (3).	<i>Lophonectes gallus.</i>
<i>Cynthia pallida.</i>	<i>Megalops cyprinoides.</i>
<i>Goodsiria coccinea</i> (3).	<i>Merluccius gayi.</i>
<i>Leptoclinum albidum</i> (3).	<i>Mesoprion fulviflamma.</i>
<i>Synstyela incrustans.</i>	<i>Monacanthus setifer.</i>
<i>Acanthurus blochi</i> (3).	<i>Muræna nebulosa.</i>
<i>Anguilla mauritiana</i> (3).	" <i>richardsoni.</i>
<i>Apogon lineatus.</i>	<i>Myxine australis.</i>
<i>Balistes buniva.</i>	<i>Notothenia longipes.</i>
<i>Bdellostoma cirrhatum.</i>	<i>Ostracion cornutus.</i>
<i>Belone annulata.</i>	<i>Platax orbicularis.</i>
<i>Brachypleura novæ-zealandiæ.</i>	<i>Platycephalus insidiator</i> (4).
<i>Callionymus curvicornis.</i>	" sp. (?)
" <i>lunatus.</i>	<i>Pseudorhombus russelli.</i>
<i>Caranx ascensionis.</i>	<i>Pseudoscarus nuchipunctatus.</i>
" <i>crumenophthalmus</i> (3).	<i>Pterois lunulata.</i>
" <i>speciosus.</i>	<i>Raja brachyura.</i>
<i>Carcharias obscurus.</i>	<i>Rhomboichthys spilurus.</i>
<i>Conger marginatus.</i>	<i>Salilota australis</i> (3).
<i>Cyprinus auratus.</i>	<i>Sebastes percooides.</i>
<i>Dules marginatus</i> (3).	<i>Serranus diacanthus</i> (3)
<i>Echeneis naucrates.</i>	<i>Solea ovata.</i>
<i>Eleotris fusca</i> (3).	<i>Sphærodon grandoculis.</i>
" <i>macrolepidota.</i>	<i>Tetrauroge longispinis.</i>
<i>Equula fasciata.</i>	<i>Tetrodon immaculatus.</i>
<i>Gerres oyena.</i>	<i>Therapon servus.</i>
<i>Glyphtodon saxatilis.</i>	<i>Trachurus trachurus.</i>
<i>Gobius albopunctatus.</i>	<i>Trygon pastinaca.</i>
<i>Haplochiton zebra</i> (3).	<i>Trypauchen vagina</i> (3).
<i>Hemiramphus commersoni.</i>	<i>Zygæna malleus.</i>
<i>Holocentrum longipinne.</i>	<i>Mesoplodon layardi.</i>

The majority of these were taken at two Stations, while 127 species were taken at three Stations, 34 species at four Stations, 19 species at five Stations, 4 species at six Stations, 1 species at seven Stations, and 1 species at nine Stations.

Of the 350 species taken in this zone less than 100 fathoms, as well as in deeper zones (those without asterisks),

285 species (or 81 per cent.) were taken in the zone between 100 and 500 fathoms.

49	" (or 14 per cent.)	"	"	"	"	500 and 1000	"
32	" (or 9 per cent.)	"	"	"	"	1000 and 1500	"
14	" (or 4 per cent.)	"	"	"	"	1500 and 2000	"
7	" (or 2 per cent.)	"	"	"	"	2000 and 2500	"
5	" (or 1 per cent.)	"	"	"	"	beyond 2500 fathoms.	