

<i>Placostegus.</i>	MACRURA :	CEPHALOPODA :
<i>Staurocephalus.</i>	<i>Brachycarpus</i> , n.g.	<i>Rossia.</i>
<i>Terebella.</i>	<i>Hepomadus</i> , n.g.	<i>Taonius.</i>
OSTRACODA :	<i>Pandalus.</i>	POLYZOA :
<i>Paradoxostoma.</i>		<i>Aetea.</i>
AMPHIPODA :	ANOMURA :	* <i>Canda.</i>
<i>Amaryllis.</i>	<i>Anapagurus.</i>	<i>Cribrilina.</i>
<i>Anonyx.</i>	* <i>Cryptodromia.</i>	<i>Flustramorpha.</i>
<i>Euthemisto.</i>	<i>Lithodes.</i>	* <i>Hippothoa.</i>
<i>Halirages.</i>	BRACHYURA :	<i>Menipea.</i>
<i>Pardalisca.</i>	* <i>Achæus.</i>	<i>Micropora.</i>
<i>Pleustes.</i>		TUNICATA :
<i>Rhachotropis.</i>	PYCNOGONIDA :	* <i>Ascidia.</i>
ISOPODA :		<i>Bathyoncus</i> , n.g.
<i>Bathytanais</i> , n.g.	LAMELLIBRANCHIATA :	<i>Boltenia.</i>
<i>Ischnosoma.</i>	<i>Astarte.</i>	<i>Clavelina.</i>
<i>Neotanais.</i>	<i>Modiolaria.</i>	<i>Didemnum.</i>
<i>Typhlotanais.</i>	<i>Saxicava.</i>	<i>Hypobythius</i> , n.g.
CUMACEA :	<i>Yoldia.</i>	<i>Styela.</i>
<i>Diastylis.</i>	GASTEROPODA :	FISHES :
<i>Leucon.</i>	<i>Buccinum.</i>	<i>Engraulis.</i>
SCHIZOPODA :		<i>Haloporphyrus.</i>
<i>Boreomysis.</i>	POLYPLACOPHORA :	<i>Latilus.</i>
	<i>Chiton.</i>	<i>Myxine.</i>
		<i>Notacanthus.</i>
		<i>Raja.</i>
		<i>Syngnathus.</i>
		<i>Trachurus.</i>

The following 34 genera, represented in the Challenger collections by species taken to the *south* of the tropics, are recorded from other sources to the *north* of the tropics, and, as far as can be gathered from the Challenger Zoological Reports, are not known from the intervening tropical zone :—

MONAXONIDA :	TETRACTINELLIDA :	HEXACTINELLIDA :
<i>Iophon.</i>	<i>Astrella.</i>	<i>Acanthascus.</i>
	<i>Stellella.</i>	<i>Rhabdocalyptus.</i>
	<i>Stryphnus.</i>	<i>Rossella.</i>