

SCYLLIIDÆ

Scyllium canicula, Cuv. (rough hound), 1910, Stations 3, 14, 20, 39.

Pristiurus melanostomus, Bonap. (black-mouthed dogfish), 1902, Faroe-Shetland channel; 1910, Stations 1, 21.

Pristiurus murinus, Coll., 1902, Faroe-Shetland channel, 1100 to 1300 metres.

CARCHARIIDÆ

Mustelus vulgaris, Müll. and Henle (smooth hound), 1910, Station 13.

LAMNIDÆ

Carcharodon, fossil tooth, 1910, Station 48 (see Fig. 254).

Oxyrhina, fossil tooth, 1910, Station 48.




FIG. 254.

Carcharodon megalodon, Fossil Tooth. Station 48. (After Zittel.) This figure shows a *Carcharodon* tooth from Tertiary deposits; those dredged from the deep-sea deposits have never the base preserved (see Fig. 126, p. 156).

SPINACIDÆ

Centrina salviani, Risso, 1910, Station 13.

Acanthias vulgaris, Risso (dog-fish), 1902, Faroe Bank, 390 metres; Faroe-Shetland channel; 1910, Stations 1, 3, 20, 39 (see Fig. 255).




FIG. 255.

Acanthias vulgaris, Risso. (After Smitt.)