

Sub-Order—HAPLOMI

SCOPELIDÆ

Bathysaurus ferox, Günth., 1910, Stations 25, 35, 53, 95 (see Fig. 103, *a*).

Bathypterois longipes, Günth., 1910, Station 53.

Bathypterois dubius, Vaill., 1910, Stations 23, 41 (see Fig. 266).




FIG. 266.

Bathypterois dubius, Vaill. Nat. size, 17 cm.

Benthosaurus grillator, G. and B., 1910, Station 53.

Bathymicrops regis, n.g., n.sp., 1910, Station 48 (see Fig. 305).

Sub-Order—HETEROMI

HALOSAURIDÆ

Halosauropsis macrochir, Günth. (Coll.), 1910, Stations 35, 53, 88, 95 (see Fig. 103, *b*).

NOTACANTHIDÆ

Notacanthus bonapartii, Risso, 1902, Faroe-Shetland channel; Faroe Bank, 840 metres (see Fig. 267).

Polyacanthonotus sp., 1910, Stations 53, 95.




FIG. 267.

Notacanthus bonapartii, Risso. (After Goode and Bean.)

Sub-Order—CATOSTEOMI

CENTRISCIDÆ

Centriscus scolopax, L., 1910, Station 39 (see Fig. 268).