

Euphorbia hypericifolia, Linn.

Euphorbia hypericifolia, Linn., Hort. Cliff., p. 198; DC., Prodr., xv. 2, p. 23; Hook., Exot. Fl., t. 36; Griseb., Fl. Brit. W. Ind., p. 54.

FERNANDO-NORONHA. Moseley.

Very common and widely spread in the warmer parts of America.

Euphorbia pilulifera, Linn.

Euphorbia pilulifera, Linn., Amoen. Acad., iii. p. 114; DC., Prodr., xv. 2, p. 21; Jacq., Ic. Pl. Rar., iii. t. 478; Griseb., Fl. Brit. W. Ind., p. 54.

FERNANDO-NORONHA. Moseley.

Generally spread in tropical countries.

Euphorbia thymifolia, Burm.

Euphorbia thymifolia, Burm., Fl. Ind., p. 2; DC., Prodr., xv. 2, p. 47; Griseb., p. 53, sub *Euphorbia maculata*.

FERNANDO-NORONHA. Moseley.

Generally spread in tropical countries, and extending into some subtropical regions.

Phyllanthus brasiliensis, Müll.

Phyllanthus brasiliensis, Müll., Arg. in DC. Prodr., xv. 2, p. 383 (*varietas genuina*).

Phyllanthus conami, Swartz; Griseb., Fl. Brit. W. Ind., p. 33.

FERNANDO-NORONHA. Moseley.

West Indies, Venezuela, Eastern Peru, Guiana, and Brazil; also recorded by Grisebach from Yucatan.

Jatropha gossypifolia, Linn.

Jatropha gossypifolia, Linn., Sp. Pl., ed. 1, p. 1006; DC., Prodr., xv. 2, p. 1086 (*varietates*); Griseb., Fl. Brit. W. Ind., p. 36.

FERNANDO-NORONHA. Moseley.

Widely diffused in Tropical America, and occurring in West Africa.

Jatropha urens, Linn.

Jatropha urens, Linn., Sp. Pl., ed. 1, p. 1007; DC., Prodr., xv. 2, p. 1100 (*varietates*); Jacq., Hort. Vindob., i. p. 8, t. 21. ¶

FERNANDO-NORONHA. Moseley.

Southern States of North America, through Mexico, to Brazil.

In addition to the foregoing Euphorbiaceæ, Sir Joseph Hooker's manuscript list of the plants collected by Mr Darwin in Fernando-Noronha contains two species of *Acalypha* and one species of *Sapium*, which we have not seen, as, for some reason, they were not sent with the others from Cambridge.