

Caulerpa mexicana, Sonder.

Caulerpa mexicana, Sonder; Kütz., Sp. Alg., p. 496; Harv., Nereis Bor.-Am., iii. p. 16, t. 37, A.

FERNANDO-NORONHA. *Moseley*.

Gulf of Mexico; Florida Keys; Bermudas; Cape Verde Islands.

Caulerpa webbiana, Mont.

Caulerpa webbiana, Mont., Syll. Gen. et Sp. Crypt., p. 453; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Darwin*.

Canary Islands.

Halimeda opuntia, Lamour.

Halimeda opuntia, Lamour.; Kütz., Sp. Alg., p. 504; Harv., Nereis Bor.-Am., iii. p. 23, t. 40, B; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley*.

Florida, West Indies, and widely dispersed in warm seas.

ULVACEÆ.

Enteromorpha — ?

Enteromorpha — ? ex Dickie in Journ. Linn. Soc. Lond., xiv. p. 365.

FERNANDO-NORONHA. *Moseley*.

So very imperfect that the genus only can be indicated; it seems nearly allied to *Enteromorpha clathrata*, Grev.

Kallonema obscurum, n. sp.

Kallonema obscurum, n. sp., Dickie in Journ. Linn. Soc. Lond., xiv. p. 365.

Frons capillaris, undulata, pallide virescens, vage et irregulariter ramosa, e cellulis subquadratis laxè connexis formata.

FERNANDO-NORONHA. *Moseley*.

The specimens are very fragmentary, mixed with some of the following in rock-pools.

Ulva lobata, Kütz.

Ulva lobata, Kütz., Sp. Alg.; Dickie in Journ. Linn. Soc. Lond., xiv. p. 365.

FERNANDO-NORONHA. *Moseley*.

Chili; Cape of Good Hope; Mauritius; Ceylon; Macao; Singapore.