

CRYPTONEMIACEÆ.

Grateloupia flicina, K. A. Agardh.

Grateloupia flicina, K. A. Agardh, Sp. Alg., i. p. 223; Kütz., Sp. Alg., p. 730; Dickie in Journ. Linn. Soc. Lond., xiii. p. 180; Melliss, St Hel., p. 381.

ST HELENA.—Rather common—*Melliss*.

Both the ordinary form and the var. *ramentacea* occur.

Generally diffused.

CERAMIACEÆ.

Centroceras clavulatum, J. G. Agardh.

Centroceras clavulatum, J. G. Agardh, Sp. Alg., ii. p. 148; Dickie in Journ. Linn. Soc. Lond., xiii. p. 180; Melliss, St Hel., p. 381.

ST HELENA.—Found adhering to *Corallina carinata*—*Melliss*.

Generally diffused.

Griffithsia setacea, K. A. Agardh.

Griffithsia setacea, K. A. Agardh, Sp. Alg., ii. p. 129; Kütz., Sp. Alg., p. 660; Dickie in Journ. Linn. Soc. Lond., xiii. p. 180; Melliss, St Hel., p. 381.

ST HELENA.—Adhering to *Laurencia cruciata*—*Melliss*.

Eastern North Atlantic; Mediterranean; New Zealand; Tasmania.

CHLOROSPERMEÆ.

SIPHONACEÆ.

Codium tomentosum, K. A. Agardh.

Codium tomentosum, K. A. Agardh, Sp. Alg., i. p. 452; Dickie in Journ. Linn. Soc. Lond., xiii. p. 180; Melliss, St Hel., p. 381.

ST HELENA.—In pools on the windward coast; very rare—*Melliss*.

Generally diffused.

VALONIACEÆ.

Valonia verticillata, Kütz.

Valonia verticillata, Kütz., Sp. Alg., p. 508? Dickie in Journ. Linn. Soc. Lond., xiii. p. 181; Melliss, St Hel., p. 381.

ST HELENA.—Rare in pools—*Melliss*.