

Mr Moseley collected the typical plant in the Cape Verde Islands; Kützing states that it is found "ad insulam Sanctæ Crucis," which by a slip is given as Vera Cruz by Dickie, whom Melliss follows.

#### ULVACEÆ.

##### *Ulva latissima*, Kütz.

*Ulva latissima*, Kütz.; Dickie in Journ. Linn. Soc. Lond., xiii. p. 181; Melliss, St Hel., p. 380.

ST HELENA.—Very common on the shores of the island—*Melliss*.

Very widely spread.

##### *Ulva parvula*, Kütz.

*Ulva parvula*, Kütz., Sp. Alg., p. 474?; Melliss, St Hel., p. 380.

ST HELENA.—Rare; adhering to *Laurencia cruciata*—*Melliss*.

The original *Ulva parvula* is from the Mediterranean. Dickie does not include this in his enumeration.

##### *Enteromorpha*, sp. ?

*Enteromorpha*, sp. ? Dickie in Journ. Linn. Soc. Lond., xiii. p. 181; Melliss, St Hel., p. 380.

ST HELENA.—Adhering to *Laurencia cruciata*—*Melliss*.

"Structure evidently the same as that of *Enteromorpha percursa*, J. G. Agardh, but too imperfect for identification."—*Dickie*.