

GRAMINEÆ.

Poa cookii, Hook. f.

Poa cookii, Hook. f. in Phil. Trans. Roy. Soc. Lond., clxviii. p. 22.

Festuca cookii, Hook. f. Fl. Antarct., p. 382, t. 139; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389 (varietas).

HEARD ISLAND. *Moseley*.

The typical plant is found in Kerguelen and Marion Islands.

[*Agrostis magellanica* (*Agrostis antarctica*) is recorded from Heard Island in the Botany of Kerguelen Island, p. 21, but this is an error.]

CRYPTOGAMÆ.

MUSCI.

Ceratodon purpureus, Brid.

Ceratodon purpureus, Brid.; Hook. f. Fl. Antarct., p. 131; Mitt. in Journ. Linn. Soc. Lond., xv. p. 73.

HEARD ISLAND.—In a small, starved, and blackened state, barren—*Moseley*.

One of the most generally diffused of all mosses.

Grimmia (*Schistidium*) *insularis*, Mitt. (Plate LIII.)

Grimmia (*Schistidium*) *insularis*, Mitt. in Journ. Linn. Soc. Lond., xv. p. 73.

Monoica. Pulvinato-cæspitosa. Folia patentia, siccitate subfalcatis curvata, lanceolata, subulato-angustata, superne carinata, integerrima, nervo crassiusculo percursa, cellulis superioribus rotundis obscuris, basalibus ad nervum angustis elongatis, ad angulos oblongis rectangularibus hyalinis; perichætialia erecta, convoluta, ovali-elliptica, acuminata, cellulis basalibus oblongis angulatis, superioribus parvis oblongis rotundisque intermixtis. Theca immersa, subrotunda, ore amplo, operculo conico-acuminato, peristomio dentibus brevioribus.

HEARD ISLAND. *Moseley*.

Grimmiæ maritimæ statura habituque simillima sed foliis apice angustiore acuminatis et basi cellulis diversiformibus areolatis margine nullibi recurvis.

Bartramia robusta, Hook. f. et Wils.

Bartramia robusta, Hook. f. et Wils. Fl. Antarct., p. 133, t. 59, fig. 4; Mitt. in Journ. Linn. Soc. Lond., xv. p. 73.

HEARD ISLAND.—A small state of what appears to be this species is intermixed with *Ceratodon purpureus*—*Moseley*.

Also in Lord Auckland and Campbell Islands.