

Gouania microcarpa, DC.

Gouania microcarpa, DC. ; Miq., Fl. Ind. Bat., i. 1, p. 650 ; Hook. f., Fl. Brit. Ind., i. p. 643 ; Seem., Fl. Vit., p. 43.

TIMOR LAUT.—Java, Malacca, and the Western Peninsula of India. *Gouania richii*, A. Gr., of the Fiji Islands, is very near this. The genus is generally diffused in the tropics, except Australia.

AMPELIDEÆ.

Vitis repens, Wight and Arnott.

Vitis repens, Wight and Arnott ; Hook. f., Fl. Brit. Ind., i. p. 646.

Cissus repens, Lam. ; Miq., Fl. Ind. Bat., i. 2, p. 605.

TIMOR LAUT.—Amboina and Java, and almost all over India. According to Miquel, the leaves are employed as an acidulant.

Vitis tenuifolia, Wight and Arnott ?

Vitis tenuifolia, Wight and Arnott ? Hook. f., Fl. Brit. Ind., i. p. 660.

TIMOR LAUT.—Nearly all over India, and also in China and Japan. Fragments of several other species of *Vitis* were collected in Arrou, Maru, Timor Laut, &c.

Leea sambucina, Willd. var. ?

Leea sambucina, Willd. var. ? Benth., Fl. Austr., i. p. 451 ; Miq., Fl. Ind. Bat., i. 2, p. 611 ; Hook. f., Fl. Brit. Ind., i. p. 666 ; Seem., Fl. Vit., p. 44.

ARROU ; DAMMAR.—This handsome shrub is generally diffused in Tropical Asia, Africa, Australia, and the Fiji Islands. Fruit a berry, as in *Vitis*. *Leea* is peculiar to the Old World.

Leea æquata, Linn.

Leea æquata, Linn., Mant., p. 124 ; C. B. Clarke in Journ. Bot., 1881, p. 163.

ARROU ; KI ; BABAR ; DAMMAR ; TIMOR LAUT.—A very common species in Eastern Tropical India and Malaya.

SAPINDACEÆ.

Erioglossum edule, Blume.

Erioglossum edule, Blume ; Miq., Fl. Ind. Bat., i. 2, p. 574 ; Benth., Fl. Austr., i. p. 454 ; Hook. f., Fl. Brit. Ind., i. p. 672.

TIMOR LAUT.—Northern Australia, and generally spread in Tropical Asia. Miquel states that the fruit of this large tree is eaten in the Archipelago. *Erioglossum* is a small genus limited to the tropics of the Old World.