

There are imperfect specimens of another species in Mr Moseley's Arrou collection. The genus *Diospyros* comprises upwards of 150 species spread over all tropical countries, and a few extending into north temperate regions in Japan, South Europe, and North America.

OLEACEÆ.

Jasminum sambac, Ait.

Jasminum sambac, Ait. ; Miq., Fl. Ind. Bat., ii. p. 542 ; Hook. f., Fl. Brit. Ind., iii. p. 591.

ARROU.—Widely dispersed in Tropical Asia, where, however, as elsewhere in the tropics, it is commonly cultivated. There are nearly a hundred species of the genus, spread over the warmer parts of the Old World ; one species reaching the south of Europe.

Jasminum didymum, Forst.

Jasminum didymum, Forst. ; Benth., Fl. Austr., iv. p. 294 ; Seem., Fl. Vit., p. 154 ; Nadeaud, Enum. Pl. Tahiti, p. 55.

Jasminum parviflorum, Dcne., Herb. Timor. Descr., p. 77 ; Miq., Fl. Ind. Bat., ii. p. 531.

TIMOR LAUT.—Java to Northern Australia ; also in New Caledonia, Tahiti, and the Fijis.

Jasminum lancifolium, Dcne. ?

Jasminum lancifolium, Dcne. ? Herb. Timor. Descr., p. 76 ; Miq., Fl. Ind. Bat., ii. p. 539.

KI.—Only seen elsewhere from Timor.

Linociera ramiflora, DC.

Linociera ramiflora, DC., Prodr., viii. p. 297.

Chionanthus ramiflora, Roxb., Fl. Ind., i. p. 107 ; Benth., Fl. Austr., iv. p. 301 ; Miq., Fl. Ind. Bat., ii. p. 551.

KI.—Tenasserim and the Philippines, to North-east Australia. Clarke (in Hook. f., Fl. Brit. Ind., iii. p. 611) makes this a variety of *Linociera macrophylla*, Wall., thus extending its range northward to the Himalayas. *Linociera* numbers nearly forty species, inhabiting the tropical regions of both hemispheres.

APOCYNACEÆ.

Rauwolfia amsoniæfolia, A. DC.

Rauwolfia amsoniæfolia, A. DC., Prodr., viii. p. 338.

KI ; TIMOR LAUT ; MARU.—Philippines, and probably some of the intermediate islands. *Rauwolfia* numbers nearly forty species, which inhabit Tropical America, Asia, Africa, and South Africa.