

Eranthemum variabile, R. Br.

Eranthemum variabile, R. Br., Prodr. Fl. N. Holl., p. 477; Benth., Fl. Austr., iv. p. 555.

TIMOR LAUT.—Common in Queensland and New South Wales, but this appears to be the first discovery of the species outside of Australia. The plant figured under this name in Paxton's Magazine of Botany, xiii. p. 75, is a distinct species. The genus is generally spread over the tropics.

Eranthemum album, Nees.

Eranthemum album, Nees; Hook. f., Fl. Brit. Ind., iv. p. 498.

Eranthemum crenulatum, Wall.; Bot. Reg., t. 879; DC., Prodr., xi. p. 453; Miq., Fl. Ind. Bat., ii. p. 836.

ARROU ; KI ; DAMMAR.—Andamans, and Malayan Peninsula and Archipelago.

Lepidagathis hyalina, Nees var. ?

Lepidagathis hyalina, Nees var. ? Miq., Fl. Ind. Bat., ii. p. 815.

TIMOR LAUT ; DAMMAR ; LETTI.—Common in India, and extending from China through the Archipelago. About fifty species of *Lepidagathis* are known, two of which are American, and the rest Old World, chiefly Asiatic.

Lepidagathis javanica, Blume.

Lepidagathis javanica, Blume; Miq., Fl. Ind. Bat., ii. p. 813.

ARROU ; KI.—Java to Timor.

Justicia gendarussa, Linn.

Justicia gendarussa, Linn. f., Suppl., p. 85; Rumph., Herb. Amb., iv. p. 70, t. 28.

Gendarussa vulgaris, Nees in DC. Prodr., xi. p. 410; Miq., Fl. Ind. Bat., ii. p. 831.

TIMOR LAUT ; BABAR.—Southern India, China, Formosa, and the Philippines through the Archipelago, but not recorded from Australia. It is also found in Madagascar, the Seychelles, Bourbon, and Mauritius; and it exists as an introduced plant in South America. *Justicia* is a genus of more than 100 species, spread over the warm regions of both hemispheres.

Graptophyllum hortense, Nees.

Graptophyllum hortense, Nees; Wall., Pl. As. Rar., iii. p. 285; DC., Prodr., xi. p. 328; Miq., Fl. Ind. Bat., ii. p. 824; T. Anders. in Journ. Linn. Soc. Lond., ix. p. 523; Dcne., Herb. Timor. Descr., p. 54; Rumph., Herb. Amb., iv. t. 30.

Justicia picta, Vahl, Symb., ii. p. 14; Bot. Reg., t. 1227; Bot. Mag., t. 1870.

ARROU ; TIMOR LAUT.—This is commonly cultivated throughout Tropical Asia, as well as in other parts of the world, on account of its showy flowers, and prettily variegated