

second species of the same genus; both from the Arrou Islands. There is also what appears to be an *Aglaonema* from the same islands, and a large fruit-bearing specimen of an Aroid from Ki.

NAIADACEÆ.

Zostera nana, Roth.

Zostera nana, Roth.; Kunth, Enum. Pl., iii. p. 19; Benth., Fl. Austr., vii. p. 176.

Zostera muelleri, Irmisch; Aschers. in Linnæa, xxxv. p. 168.

ARROU.—Coasts of Southern Australia, Tasmania, and New Zealand, as well as in the northern hemisphere in the Old World.

CYPERACEÆ.

Cyperus ferax, A. Rich.

Cyperus ferax, A. Rich.; Benth., Fl. Austr., vii. p. 286; Clarke in Journ. Linn. Soc. Lond., xx. p. 295, et xxi. p. 191.

TIMOR LAUT.—Widely spread over the tropics, both in America and the Old World.

Cyperus pennatus, Lam.

Cyperus pennatus, Lam.; Benth., Fl. Austr., vii. p. 284; Miq., Fl. Ind. Bat., iii. p. 281; Seem., Fl. Vit., p. 319; Nadeaud, Enum. Pl. Tahiti, p. 33.

TIMOR LAUT.—Widely spread in Tropical Asia and Polynesia; also in Australia and the Seychelles. It is a sea-shore plant.

GRAMINEÆ.

Panicum sanguinale, Linn.

Panicum sanguinale, Linn.; Benth., Fl. Austr., vii. p. 469; Seem., Fl. Vit., p. 325; Nadeaud, Enum. Pl. Tahiti, p. 31.

Digitaria sanguinalis, Scop.; Miq., Fl. Ind. Bat., iii. p. 437.

TIMOR LAUT.—This grass is found almost everywhere in tropical and subtropical countries.

Setaria macrostachya, H.B.K.

Setaria macrostachya, H.B.K., Nov. Gen. et Sp., i. p. 110; Benth., Fl. Austr., vii. p. 493.

Setaria italica, Miq., Fl. Ind. Bat., iii. p. 467, non Beauv.?

TIMOR LAUT.—Common in many parts of Tropical America; also occurring in India, but less commonly, according to Bentham, and in Eastern Australia.

Setaria verticillata, Beauv.

Setaria verticillata, Beauv., Agrost., p. 51; Kunth, Enum. Pl., i. p. 152; Benth., Fl. Austr., vii. p. 494.

TIMOR LAUT.—Almost universally dispersed, as a weed of cultivation, in warm and temperate countries.