

Polyporus (Mesopus) xanthopus, Fries.

Polyporus (Mesopus) xanthopus, Fries, Obs., ii. p. 285; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Spread over all tropical countries.

Polyporus (Pleuropus) sanguineus, Fries.

Polyporus (Pleuropus) sanguineus, Fries, Ep., p. 444; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 45.

KI.—This is spread all over the tropics.

Polyporus (Pleuropus) affinis, Nees.

Polyporus (Pleuropus) affinis, Nees in Fries, Ep., p. 445; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Malayan Archipelago; Polynesia; Brazil.

Polyporus (Pleuropus) cochlear, Nees.

Polyporus (Pleuropus) cochlear, Nees in Nov. Act. Nat. Cur., xiii. t. 6; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Java.

Polyporus (Pleuropus) luteus, Nees.

Polyporus (Pleuropus) luteus, Nees in Fries, Ep., p. 445; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Nicobar Islands; Australia; Cuba; Brazil.

Polyporus (Pleuropus) meleagris, Berkl.

Polyporus (Pleuropus) meleagris, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

Pileo flabelliformi stipite brevissimo affixo zonato radiato lineato brunneo; margine lobato acutissimo quandoque striato; hymenio e pallido umbrino; poris angulatis hic illic sinuatis, dissepimentis rigidiusculis.

Pileus 2–3 inches across, $1\frac{1}{2}$ inches long; pores $\frac{1}{16}$ inch in diameter. Sometimes infundibuliform. There is a variety (a single specimen only) with a longer stem, which is slightly laccate.

ARROU.

Polyporus (Anodermei) armeniacus, Berkl.

Polyporus (Anodermei) armeniacus, Berkl. in Hook. Kew Journ. Bot., 1856, p. 197; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Brazil.