

Hexagona similis, Berkl.

Hexagona similis, Berkl. in Hook. Lond. Journ. Bot., 1846, p. 4; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43 (*ubi errore affinis, fide Cooke*).

ARROU.—Ceylon ; Australia.

Favolus saccharinus, Berkl.

Favolus saccharinus, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Pileo flabelliiformi sessili subtiliter velutino glabrescente ; poris sinuatis flexuosis.

Pileus $\frac{1}{2}$ to 1 inch across, at first very minutely velvety, so as to have a saccharine appearance ; pores arranged in lines proceeding from a common base.

ARROU.

Favolus brasiliensis, Fries.

Favolus brasiliensis, Fries, Ep., p. 498 ; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

ARROU.—North India ; Ceylon ; Southern United States ; Cuba ; Surinam ; Brazil.

Laschia longipes, Berkl.

Laschia longipes, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Pileo tenui umbilicato pallide aurantiaco, margine lobato ; stipite elongato sursum dilatato fissili ; poris brevibus angustis sinuatis radiantibus.

About 1 inch across ; stem 2 inches high, not $\frac{1}{2}$ inch thick. Specimen bad, but very distinct.

ARROU.

Dædalea indurata, Berkl.

Dædalea indurata, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

Pileo albido, disco parvo affixo radiata-rugoso marginem versus zonato ; hymenio ochroleuco ; poris minutis sinuositis.

Pileus $1\frac{1}{2}$ inches across, 1 long ; pores $\frac{1}{60}$ inch in diameter. A very distinct species, of which there is a single specimen only.

ARROU.

Irpea maximus, Mont.

Irpea maximus, Mont., Syll., p. 174 ; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44.

ARROU.—Cuba ; San Domingo.