

ENUMERATION OF THE PLANTS.¹

DICOTYLEDONES.—POLYPETALÆ.

MENISPERMACEÆ.

Stephania hernandiæfolia, Walp.

Stephania hernandiæfolia, Walp.; Benth., Fl. Austr., i. p. 57; Miq., Fl. Ind. Bat., i. 2, p. 83; Hook. f., Fl. Brit. Ind., i. p. 103.

ADMIRALTY ISLANDS.—Spread nearly all over Tropical Asia and Africa, and extending in Asia northward to Japan. In Australia it ranges from Cape York to Victoria, and there are specimens in the Kew Herbarium from Tahiti and Samoa. As circumscribed by Bentham, this includes *Stephania gaudichaudi*, A. Gr. (Bot. U.S. Expl. Exped., i. p. 37), and several other named forms. *Stephania forsteri*, A. Gr., *loc. cit.*, p. 36 (*Cocculus forsteri*, DC.), is also apparently the same species. Seemann does not mention this plant in his Flora Vitiensis.

GUTTIFERÆ.

Ochrocarpus ovalifolius, T. Anders.

Ochrocarpus ovalifolius, T. Anders., MSS. in Herb. Kew.

Calysaccion ovalifolium, Chois.; Miq., Fl. Ind. Bat., i. 2, p. 512.

Calophyllum excelsum, Zoll. in Nat. en Geneesk. Archief Néerl. Ind., ii. p. 558, *vide* Miq.

ADMIRALTY ISLANDS.—Malayan Archipelago.

Calophyllum inophyllum, Linn.

Calophyllum inophyllum, Linn.; Seem., Fl. Vit., p. 12; Nadeaud, Enum. Pl. Tahiti, p. 69; Jouan in Mém. Soc. Sci. Nat. Cherbourg, xi., 1865, p. 101; Mann in Proc. Amer. Acad., vii. p. 156.

ADMIRALTY ISLANDS.—A conspicuous seaside tree in Tropical Asia, Africa, Australia, and Polynesia.

¹ Full references and details of the distribution of those species which occur in the preceding enumeration of plants from the South-eastern Moluccas are not repeated here; the references being limited to the Polynesian literature, with an outline of the distribution.